

GRAPHICAL MODELLING OF MULTIVARIATE TIME SERIES

MICHAEL EICHLER

*Department of Quantitative Economics, University of Maastricht
P.O. Box 616, 6200 MD Maastricht, The Netherlands*

February 2, 2008

ABSTRACT. We introduce graphical time series models for the analysis of dynamic relationships among variables in multivariate time series. The modelling approach is based on the notion of strong Granger causality and can be applied to time series with non-linear dependencies. The models are derived from ordinary time series models by imposing constraints that are encoded by mixed graphs. In these graphs each component series is represented by a single vertex and directed edges indicate possible Granger-causal relationships between variables while undirected edges are used to map the contemporaneous dependence structure. We introduce various notions of Granger-causal Markov properties and discuss the relationships among them and to other Markov properties that can be applied in this context.

Keywords: Graphical models, multivariate time series, Granger causality, global Markov property

1. INTRODUCTION

Graphical models have become an important tool for the statistical analysis of complex multivariate data sets, which are now increasingly available in many scientific fields. The key feature of these models is to merge the probabilistic concept of conditional independence with graph theory by representing possible dependencies among the variables of a multivariate distribution in a graph. This has led to simple graphical criteria for identifying the conditional independence relations that are implied by a model associated with a given graph. Further important advantages of the graphical modelling approach are statistical efficiency due to parsimonious parameterizations of the joint distribution of the variables and the visualization of complex dependence structures, which allows an intuitive understanding of the interrelations among the variables and, thus, facilitates the communication of statistical results. For an introduction to graphical models we refer to the monographs by Whittaker (1990), Edwards (2000), and Cox and Wermuth (1996); a mathematically more rigorous treatment can be found in Lauritzen (1996).

While graphical models originally have been developed for variables that are sampled with independent replications, they have been applied more recently also to the analysis of time dependent data. Some first general remarks concerning the potential

The paper was written while the author was working at the Institut für Angewandte Mathematik, Universität Heidelberg, Germany.

E-mail address: m.eichler@ke.unimaas.nl (M. Eichler).

use of graphical models in time series analysis can be found in Brillinger (1996); since then there has been an increasing interest in the use of graphical modelling techniques for analyzing multivariate time series (e.g., Stanghellini and Whittaker 1999, Dahlhaus 2000, Reale and Tunnicliffe Wilson 2001, Dahlhaus and Eichler 2003, Oxley et al. 2004, Moneta and Spirtes 2005, Eichler 2006a,b). However, all these works have been restricted to the analysis of linear interdependencies among the variables whereas the recent trend in time series analysis has shifted towards non-linear parametric and non-parametric models (e.g., Tong 1993, Rothman 1999, Fan and Yao 2003). Moreover, in most of these approaches, the variables at different time points are represented by separate nodes, which leads to graphs with theoretically infinitely many vertices for which no rigorous theory exists so far.

In this paper, we present a general approach for graphical modelling of multivariate stationary time series, which is based on simple graphical representations of the dynamic dependencies of a process. To this end, we utilize the concept of strong Granger causality (e.g., Florens and Mouchart 1982), which is formulated in terms of conditional independencies and, thus, can be applied to model arbitrary non-linear relationships among the variables. The concept of Granger causality originally has been introduced by Granger (1969) and is commonly used for studying dynamic relationships among the variables in multivariate time series.

For the graphical representations, we consider mixed graphs in which each variable as a complete time series is represented by a single vertex and directed edges indicate possible Granger-causal relationships among the variables while undirected edges are used to map the contemporaneous dependence structure. We note that similar graphs have been used in Eichler (2006a) as path diagrams for the autoregressive structure of weakly stationary processes. Formally, the graphical encoding of the dynamic structure of a time series is achieved by a new type of Markov properties, which we call Granger-causal Markov properties. We introduce various levels, namely the pairwise, the local, the block-recursive, and the global Granger-causal Markov property, and discuss the relationships among them. In particular, we give sufficient conditions under which the various Granger-causal Markov properties are equivalent; such conditions allow formulating models based on a simple Markov property while interpreting the associated graph by use of the global Granger-causal Markov property.

The paper is organized as follows. In Section 2, we introduce the concepts of Granger-causal Markov properties and graphical time series models; some examples of graphical time series models are presented in Section 3. In Section 4, we discuss global Markov properties, which relate certain separation properties of the graph to conditional independence or Granger noncausality relations among the variables of the process. Finally in Section 5, we compare the presented graphical modelling approach with other approaches in the literature and discuss possible extensions. The proofs are technical and put into the appendix.

2. GRAPHICAL TIME SERIES MODELS

In graphical modelling, the focus is on multivariate statistical models for which the possible dependencies between the studied variables can be represented by a graph. In multivariate time series analysis, statistical models for a time series $X_V =$

$(X_V(t))_{t \in \mathbb{Z}}$ are usually specified in terms of the conditional distribution of $X_V(t+1)$ given its past $\mathbf{X}_V(t) = (X_V(s))_{s \leq t}$ in order to study the dynamic relationships over time among the series. Thus, a time series model may be described formally as a family of probability kernels P from $\mathbb{R}^{V \times \mathbb{N}}$ to $\mathbb{R}^V$, and we write $X_V \sim P$ if P is a version of the conditional probability of $X_V(t+1)$ given $\mathbf{X}_V(t)$.

For modelling specific dependence structures, we utilize the concept of Granger (non-)causality, which has been introduced by Granger (1969) and has proved to be particularly useful for studying dynamic relationships in multivariate time series. This probabilistic concept of noncausality from a process X_a to another process X_b is based on studying whether at time t the next value of X_b can be better predicted by using the entire information up to time t than by using the same information apart from the former series X_a . In practice, not all relevant variables may be available and, thus, the notion of Granger causality clearly depends on the used information set. In the sequel, we use the concept of strong Granger noncausality (e.g., Florens and Mouchart 1982), which is defined in terms of conditional independence and σ -algebras and, thus, can be used also for non-linear time series models.

Let $X_V = (X_V(t))_{t \in \mathbb{Z}}$ with $X_V(t) = (X_v(t))'_{v \in V} \in \mathbb{R}^V$ be a multivariate stationary stochastic process on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. For $A \subseteq V$, we denote by $X_A = (X_A(t))_{t \in \mathbb{Z}}$ the multivariate subprocess with components X_a , $a \in A$. The information provided by the past and present values of X_A at time $t \in \mathbb{Z}$ can be represented by the sub- σ -algebra $\mathcal{X}_A(t)$ of $\mathcal{F}$ that is generated by $\mathbf{X}_A(t) = (X_A(s))_{s \leq t}$. We write $\mathcal{X}_A = (\mathcal{X}_A(t), t \in \mathbb{Z})$ for the filtration induced by X_A . This leads to the following definition of (strong) Granger noncausality in multivariate time series.

Definition 2.1. Let A and B be disjoint subsets of V .

- (i) X_A is *strongly Granger-noncausal* for X_B with respect to the filtration $\mathcal{X}_V$ if

$$\mathcal{X}_B(t+1) \perp\!\!\!\perp \mathcal{X}_A(t) \mid \mathcal{X}_{V \setminus A}(t)$$

for all $t \in \mathbb{Z}$. This will be denoted by $X_A \nrightarrow X_B [\mathcal{X}_V]$.

- (ii) X_A and X_B are *contemporaneously conditionally independent* with respect to the filtration $\mathcal{X}_V$ if

$$\mathcal{X}_A(t+1) \perp\!\!\!\perp \mathcal{X}_B(t+1) \mid \mathcal{X}_V(t) \vee \mathcal{X}_{V \setminus (A \cup B)}(t+1)$$

for all $t \in \mathbb{Z}$. This will be denoted by $X_A \approx X_B [\mathcal{X}_V]$.

In the following, we will speak simply of Granger (non-)causality in the sense of the above definition.

Intuitively, the dynamic relationships of a stationary multivariate time series X_V can be visualized by a mixed graph $G = (V, E)$ in which each vertex $v \in V$ represents one component X_v and two vertices a and b are joined by a directed edge $a \rightarrow b$ whenever X_a is Granger-causal for X_b or by an undirected edge $a - b$ whenever X_a and X_b are contemporaneously conditionally dependent. Conversely, for formulating models with specific dynamic dependencies, a mixed graph G can be associated with a set of Granger noncausality and contemporaneous conditional independence constraints that are imposed on a time series model for X_V . Such a set of conditional independence relations encoded by a graph G is generally known




FIGURE 2.1. Encoding of relations $X_A \nrightarrow X_B \mid \mathcal{X}_X$ by the (a) pairwise, (b) local, and (c) block-recursive Granger-causal Markov property (A and B are indicated by grey and black nodes, respectively).

as Markov property with respect to G . In the context of multivariate time series, graphs may encode different types of conditional independence relations, and we therefore speak of Granger-causal Markov properties when dealing with Granger noncausality and contemporaneous conditional independence relations. In the following definition, $\text{pa}(a) = \{v \in V \mid v \rightarrow a \in E\}$ denotes the set of parents of a vertex a , while $\text{ne}(a) = \{v \in V \mid v - a \in E\}$ is the set of neighbours of a ; furthermore, for $A \subseteq V$, we define $\text{pa}(A) = \cup_{a \in A} \text{pa}(a) \setminus A$ and $\text{ne}(A) = \cup_{a \in A} \text{ne}(a) \setminus A$.

Definition 2.2 (Granger-causal Markov properties). Let $G = (V, E)$ be a mixed graph. Then the stochastic process X_V satisfies

- (PC) the *pairwise Granger-causal Markov property* with respect to G if for all $a, b \in V$ with $a \neq b$
 - (i) $a \rightarrow b \notin E \Rightarrow X_a \nrightarrow X_b \mid \mathcal{X}_V$,
 - (ii) $a - b \notin E \Rightarrow X_a \approx X_b \mid \mathcal{X}_V$;
- (LC) the *local Granger-causal Markov property* with respect to G if for all $a \in V$
 - (i) $X_{V \setminus (\text{pa}(a) \cup \{a\})} \nrightarrow X_a \mid \mathcal{X}_V$,
 - (ii) $X_{V \setminus (\text{ne}(a) \cup \{a\})} \approx X_a \mid \mathcal{X}_V$;
- (BC) the *block-recursive Granger-causal Markov property* with respect to G if for all subsets A of V
 - (i) $X_{V \setminus (\text{pa}(A) \cup A)} \nrightarrow X_A \mid \mathcal{X}_V$,
 - (ii) $X_{V \setminus (\text{ne}(A) \cup A)} \approx X_A \mid \mathcal{X}_V$.

Similarly, if P is a probability kernel from $\mathbb{R}^{V \times \mathbb{N}}$ to $\mathbb{R}^V$, we say that P satisfies the pairwise, the local, or the block-recursive Granger-causal Markov property with respect to a graph G whenever the same is true for every stationary process X_V with $X_V \sim P$.

Example 2.3. To illustrate the various Granger-causal Markov properties, we consider the graph G in Figure 2.1. Suppose that a stationary process X_V satisfies the pairwise Granger-causal Markov property with respect to this graph G . Then the absence of the edge $1 \rightarrow 4$ in G implies that X_1 is Granger-noncausal for X_4 with respect to $\mathcal{X}_V$. Next, in the case of the local Granger-causal Markov property, we find that the bivariate subprocess $X_{\{1,2\}}$ is Granger-noncausal for X_4 with respect to $\mathcal{X}_V$ since vertex 4 has parents 3 and 5. Similarly, if X_V obeys the block-recursive Granger-causal Markov property, the graph encodes that $X_{\{1,2\}}$ is Granger-noncausal for $X_{\{4,5\}}$ with respect to $\mathcal{X}_V$ since $\text{pa}(4, 5) = \{3\}$.

The block-recursive Granger-causal Markov property obviously implies the other two Granger-causal Markov properties and, thus, is the strongest of the three Markov

properties; similarly, the pairwise Granger-causal Markov property clearly is the weakest of the three properties. The question arises whether and under which conditions the three Granger-causal Markov properties are equivalent. In the case of random vectors $(Y_v)_{v \in V}$ with values in $\mathbb{R}^V$, the various levels of Markov properties for graphical interaction models are equivalent if the distribution of Y_V satisfies

$$Y_A \perp\!\!\!\perp Y_B \mid Y_{C \cup D} \wedge Y_A \perp\!\!\!\perp Y_C \mid Y_{B \cup D} \Rightarrow Y_A \perp\!\!\!\perp Y_{B \cup C} \mid Y_D$$

for all disjoint subsets A, B, C , and D of V (Pearl and Paz 1987). A sufficient condition for this intersection property is that the joint distribution of Y_V is absolutely continuous with respect to some product measure and has a positive and continuous density (e.g., Lauritzen 1996, Prop. 3.1). The following result establishes similar conditions for the time series case.

Proposition 2.4. *Suppose that the following two conditions hold:*

- (M) $X_V = (X_V(t))_{t \in \mathbb{Z}}$ is a stationary, strongly mixing stochastic process on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$ taking values in $\mathbb{R}^V$;
- (P) the conditional distribution $\mathbb{P}^{X_V(t+1) \mid \mathbf{X}_V(t)}$, $t \in \mathbb{Z}$, has a regular version that is almost surely absolutely continuous with respect to some product measure ν on $\mathbb{R}^{|V|}$ with ν -a.e. positive and continuous density.

Then, for every $\mathcal{F}$ -measurable random variable Y and every $t \in \mathbb{Z}$,

$$Y \perp\!\!\!\perp \mathcal{X}_A(t) \mid \mathcal{X}_{B \cup C}(t) \wedge Y \perp\!\!\!\perp \mathcal{X}_B(t) \mid \mathcal{X}_{A \cup C}(t) \Leftrightarrow Y \perp\!\!\!\perp \mathcal{X}_{A \cup B}(t) \mid \mathcal{X}_C(t).$$

With this intersection property, we obtain the following relations among the three Granger-causal Markov properties.

Theorem 2.5. *Suppose that X_V satisfies conditions (M) and (P). Then the three Granger-causal Markov properties (BC), (LC), and (PC) are related by the following implications:*

$$(BC) \Rightarrow (LC) \Leftrightarrow (PC).$$

Furthermore, if X_V additionally satisfies

$$X_A \nrightarrow X_B \mid [\mathcal{X}_V] \Leftrightarrow X_A \nrightarrow X_b \mid [\mathcal{X}_V] \quad \forall b \in B, \quad (2.1)$$

then the three Granger-causal Markov properties (BC), (LC), and (PC) are equivalent.

The theorem shows that, similarly as in the case of chain graph models with the Andersson-Madigan-Perlman (AMP) Markov property (Andersson et al. 2001), the pairwise and the local Granger-causal Markov property are in general not sufficiently strong to encode all Granger-causal relationships that hold among the components of a multivariate time series with respect to full information $\mathcal{X}_V$. This suggests to specify graphical time series models in terms of the block-recursive Granger-causal Markov property.

Definition 2.6 (Graphical time series model). Let G be a mixed graph and let $\mathcal{P}_G$ be a statistical time series model given by a family of probability kernels $P \in \mathcal{P}_G$ from $\mathbb{R}^{V \times \mathbb{N}}$ to $\mathbb{R}^V$. Then $\mathcal{P}_G$ is said to be a *graphical time series model* associated with the graph G if, for all $P \in \mathcal{P}_G$, the distribution P satisfies the block-recursive Granger-causal Markov property with respect to G .

The three Granger-causal Markov properties considered so far encode only Granger noncausality relations with respect to the complete information $\mathcal{X}_V$. The discussion of phenomena such as spurious causality (e.g., Hsiao 1982, Eichler 2005a), however, requires also the consideration of Granger-causal relationships with respect to partial information sets, that is, with respect to filtrations $\mathcal{X}_S$ for subsets S of V . To this end, we introduce in Section 4 a global Granger-causal Markov property that more generally relates pathways in a graph to Granger-causal relations among the variables, and we establish, under conditions (M) and (P), its equivalence to the block-recursive Granger-causal Markov property; this shows that the block-recursive Granger-causal Markov property is indeed sufficiently rich to describe the dynamic dependence structure in multivariate time series.

Before we continue our discussion of Markov properties in Section 4, we illustrate the introduced concept of graphical time series models by a few examples.

3. EXAMPLES

In the previous section, graphical time series models have been defined in terms of the block-recursive Granger-causal Markov property. For many time series models, however, condition (2.1) in Proposition 2.5 holds, and, hence, the pairwise and the block-recursive Granger-causal Markov property are equivalent. This enables us to derive the constraints on the parameters from the pairwise Markov property.

There are no simple conditions known that are both necessary and sufficient for (2.1). The following proposition lists some sufficient conditions that cover many examples, as will be shown subsequently. A counter-example that demonstrates that the pairwise and the block-recursive Granger-causal Markov property are in general not equivalent will be provided in Example 3.5.

Proposition 3.1. *Suppose that X_V satisfies conditions (M) and (P) and one of the following conditions:*

- (i) X_V is a Gaussian process;
- (ii) $X_v(t+1)$, $v \in V$, are mutually contemporaneously independent, that is, the joint conditional distribution factorizes as

$$\mathbb{P}^{X_V(t+1) | \mathbf{X}_V(t)} = \otimes_{v \in V} \mathbb{P}^{X_v(t+1) | \mathbf{X}_V(t)} \quad \forall t \in \mathbb{Z};$$

- (iii) $X_V(t+1)$ depends on its past only in its conditional mean, that is,

$$X(t+1) - \mathbb{E}[X(t+1) | \mathcal{X}_V(t)] \perp\!\!\!\perp \mathcal{X}_V(t) \quad \forall t \in \mathbb{Z}.$$

Then the three Granger-causal Markov properties (BC), (LC), and (PC) are equivalent.

Example 3.2 (Vector autoregressive processes). Let X_V be a stationary vector autoregressive process of order p ,

$$X_V(t) = \sum_{u=1}^p \Phi(u) X_V(t-u) + \varepsilon(t), \quad \varepsilon(t) \stackrel{\text{iid}}{\sim} \mathcal{N}(0, \Sigma), \quad (3.1)$$

where $\Phi(u)$ are $V \times V$ matrices and the variance matrix Σ is non-singular. Then X_A is Granger-noncausal for X_B with respect to $\mathcal{X}_V$ if and only if the corresponding entries $\Phi_{BA}(u)$ vanish for all $u = 1, \dots, p$ (e.g., Boudjellaba et al. 1992). Furthermore, X_A and X_B are contemporaneously conditionally independent if and only if

the corresponding error components $\varepsilon_A(t)$ and $\varepsilon_B(t)$ are conditionally independent given all remaining components $\varepsilon_{V \setminus (A \cup B)}(t)$. For Gaussian errors, conditional independencies are given by zeros in the concentration matrix $K = \Sigma^{-1}$ (e.g., Lauritzen 1996). Consequently, the process X_V satisfies the pairwise—and by Proposition 3.1(i) also the block-recursive—Granger-causal Markov property with respect to a graph $G = (V, E)$ if the following two conditions hold:

- (i) $a \rightarrow b \notin E \Rightarrow \Phi_{ba}(u) = 0 \quad \forall u = 1, \dots, p$;
- (ii) $a - b \notin E \Rightarrow K_{ab} = K_{ba} = 0$.

Thus, the graphical vector autoregressive model of order p associated with the graph G , denoted by $\text{VAR}(p, G)$, is given by the set of all stationary $\text{VAR}(p)$ processes whose parameters are constrained to zero according to the conditions (i) and (ii).

Example 3.3 (Nonparametric additive models). More generally, we may also consider nonlinear autoregressive time series models. For example, let X_V be a strongly mixing stationary process given by

$$X_b(t) = \sum_{a \in V} \sum_{u=1}^p m_{ba}^{(u)}(X_a(t-u)) + \varepsilon_b(t), \quad b \in V, t \in \mathbb{Z},$$

where $m_{ba}^{(u)}$ are measurable real-valued functions and the errors $\varepsilon_V(t)$ are independent and identically distributed. Furthermore, we assume that the distribution of the errors $\varepsilon_V(t)$ has a positive and continuous density f_V on $\mathbb{R}^V$. Then a nonparametric additive model of order p is given by the set $\mathcal{P} = \{P_{m,f}\}$ of regular conditional distributions $P_{m,f}$ of such processes X_V . For a mixed graph $G = (V, E)$, we now define $\mathcal{P}_G$ as the subset of all $P_{m,f} \in \mathcal{P}$ such that

- (i) $m_{ba}^{(u)}(\cdot)$, $u = 1, \dots, p$, are constant whenever $a \rightarrow b \notin E$ and
- (ii) f_V factorizes as $f_V(z_V) = g(z_{V \setminus \{b\}}) h(z_{V \setminus \{a\}})$ whenever $a - b \notin E$.

The second condition implies $\varepsilon_a(t) \perp\!\!\!\perp \varepsilon_b(t) \mid \varepsilon_{V \setminus \{a,b\}}(t)$ and ensures that the distribution of the errors $\varepsilon(t)$ obeys the so-called pairwise Markov property with respect to the undirected subgraph G^u of G (i.e., the subgraph G^u obtained from G by removing all directed edges). Consequently, X_a and X_b are contemporaneously conditionally independent with respect to $\mathcal{X}_V$ whenever the corresponding vertices a and b are not joined by an undirected edge in G . Similarly, the first condition implies that, for any process with $X_V \sim P_{m,f}$, the conditional distribution of $X_b(t)$ given the past $\mathbf{X}_V(t-1)$ satisfies

$$\mathbb{P}^{X_b(t) \mid \mathbf{X}_V(t-1)} = \mathbb{P}^{X_b(t) \mid \mathbf{X}_{V \setminus \{a\}}(t-1)}$$

whenever the graph G does not contain the edge $a \rightarrow b$. It follows that any process X_V with $X_V \sim P_{m,f} \in \mathcal{P}_G$ obeys the pairwise Granger-causal Markov property with respect to G . Furthermore, since $X_V(t)$ depends on its past $\mathbf{X}_V(t-1)$ only in its conditional mean, it follows from Proposition 3.1(iii) that the pairwise and the block-recursive Granger-causal Markov properties are equivalent. Thus, $\mathcal{P}_G$ is indeed the graphical nonparametric additive model of order p associated with the graph G .

Example 3.4 (Binary time series). As an example with categorical data, we consider a binary time series model that has been used for the identification of neural interactions from neural spike train data (Brillinger 1988a,b). Suppose that

the data consist of the recorded spike trains for a set of neurons, that is, of the sequences of firing times $(\tau_{v,n})_{n \in \mathbb{N}}$ for neurons $v \in V$, and let X_v be the binary time series obtained by setting $X_v(t) = 1$ if neuron v has fired in the interval $[t, t+1)$ and $X_v(t) = 0$ otherwise. Then, the interactions between the neurons can be modelled by the conditional probabilities

$$\mathbb{P}(X_b(t) = 1 | \mathcal{X}_V(t-1)) = \Phi\left(\sum_{a \in \text{pa}(b)} U_{ba}(t) - \theta\right), \quad (3.2)$$

where $\Phi(x)$ denotes the normal cumulative function,

$$U_{ba}(t) = \sum_{u=1}^{\gamma_b(t)} g_{ba}(u) X_a(t-u) \quad (3.3)$$

measures the influence of process a on process b , and

$$\gamma_b(t) = \min \{u \in \mathbb{N} | X_b(t-u) = 1\}$$

is the time elapsed since the last event of process X_b . Furthermore, we assume that the time unit has been chosen small enough such that there are no interactions among the neurons within one time interval, and that, consequently, the joint conditional probability factorizes as

$$\mathbb{P}(X_V(t) = x_V | \mathcal{X}_V(t-1)) = \prod_{v \in V} \mathbb{P}(X_v(t) = x_v | \mathcal{X}_V(t-1))$$

for all $x_V \in \{0, 1\}^V$. Then the pairwise and the block-recursive Granger-causal Markov property are equivalent by Proposition 3.1(ii) and, thus, we can use the former for modelling dependencies between the processes. From (3.2) and (3.3), it follows that X_a is Granger-noncausal for X_b if and only if $g_{ba}(u) = 0$ for all $u \in \mathbb{N}$.

Example 3.5. In general, the pairwise and the block-recursive Granger-causal Markov property are not equivalent. To demonstrate this, we consider a simple stationary Markov process X_V with conditional distributions

$$X_V(t) | \mathcal{X}_V(t-1) \sim \mathcal{N}(0, \Sigma(t))$$

and conditional covariance matrix

$$\Sigma(t) = \begin{pmatrix} 1 & \rho(t) & 0 \\ \rho(t) & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

We assume that the conditional correlation between $X_1(t)$ and $X_2(t)$ given $\mathcal{X}_V(t-1)$ depends on $X_3(t-1)$ by

$$\rho(t) = \begin{cases} \rho & \text{if } |X_3(t-1)| > c \\ 0 & \text{otherwise} \end{cases}$$

for some constants ρ with $0 < |\rho| < 1$ and $c > 0$. In other words, the variables X_1 and X_2 start becoming contemporaneously dependent once the most recent value of variable X_3 exceeds a certain threshold. For this model, we find that, on the one hand, the marginal conditional distributions of $X_v(t)$ given $\mathcal{X}_V(t-1)$ are standard normal and, thus, do not depend on $\mathcal{X}_V(t-1)$. This implies that the process X_V satisfies the pairwise Granger-causal Markov property with respect to the graph (a) in Figure 3.1. On the other hand, the bivariate conditional distribution of




FIGURE 3.1. Illustration of non-equivalence of pairwise and block-recursive Granger-causal Markov properties: the process in Example 3.5 satisfies the pairwise Granger-causal Markov property with respect to the graphs in (a) and (b) whereas it satisfies the block-recursive Granger-causal Markov property only with respect to the graph in (b).

$(X_1(t), X_2(t))$ depends on the value of $X_3(t-1)$ by the conditional correlation $\rho(t)$. Hence, X_V obeys the block-recursive Granger-causal Markov property with respect to the graph (b) in Figure 3.1, but not with respect to the graph (a).

4. GLOBAL MARKOV PROPERTIES

The interpretation of graphs describing the dependence structure of graphical models in general is enhanced by global Markov properties that merge the notion of conditional independence with a purely graph theoretical concept of separation allowing one to state whether two subsets of vertices are separated by a third subset of vertices. In this section, we show that the concept of p -separation introduced by Levitz et al. (2001) for chain graph models with the AMP Markov property (Andersson et al. 2001) can be used to obtain global Markov properties in the present context of graphical time series models. Throughout this section we assume that conditions (M) and (P) in Proposition 2.4 hold.

4.1. The global AMP Markov property

We start with some further graphical terminology. Let $G = (V, E)$ be a mixed graph. Then a *path* π between two vertices a and b in G is a sequence $\pi = \langle e_1, \dots, e_n \rangle$ of edges $e_i \in E$ such that e_i is an edge between v_{i-1} and v_i for some sequence of vertices $v_0 = a, v_1, \dots, v_n = b$. The vertices a and b are the *endpoints* of the path, while $v_1, \dots, v_{n-1}$ are the *intermediate vertices* on the path. Notice that paths may be self-intersecting since we do not require that the vertices v_j are distinct.

An intermediate vertex c on a path π is said to be a *p -collider* on the path if the edges preceding and succeeding c on the path either have both an arrowhead at c or one has an arrowhead at c and the other is a line, i.e. $\rightarrow c \leftarrow$, $\rightarrow c \text{ ---}$, $\text{---} c \leftarrow$; otherwise the vertex c is said to be a *p -noncollider* on the path. A path π between vertices a and b is said to be *p -connecting* given a set S if

- (i) every p -noncollider on the path is not in S , and
- (ii) every p -collider on the path is in S ,

otherwise we say the path is *p -blocked* given S .

Definition 4.1 (p -separation). Two vertices a and b in a mixed graph G are *p -separated* given a set S if all paths between a and b are p -blocked given C . Similarly, two sets A and B in G are said to be *p -separated* given S if, for every pair $a \in A$ and $b \in B$, a and b are p -separated given S . This will be denoted by $A \bowtie_p B \mid S$.




FIGURE 4.1. Illustration of global AMP Markov property: (a) path between 1 and 4 that is p -connecting given $S \subseteq \{2, 5\}$; (b) path between 1 and 4 that is p -connecting given $S = \{2, 3\}$ (or $\{2, 3, 5\}$); (c) path between 1 and 4 that is p -connecting given $S = \{3, 5\}$ (or $\{3\}$).

We note that the above conditions for p -separation are simpler than those in Levitz et al. (2001) due to the fact that we consider the larger class of all possibly self-intersecting paths. The equivalence of the two notions of p -separation is shown in Appendix D. The following results show that the concept of p -separation can be applied to graphs encoding dynamic relationships in multivariate time series and allows reading off conditional independencies among the stochastic processes that are represented by the vertices in the graph.

Lemma 4.2. *Suppose that X_V satisfies the block-recursive Granger-causal Markov property with respect to the graph G . Then, for any disjoint subsets A , B , and S of V , we have*

$$A \bowtie_p B \mid S \Rightarrow \mathcal{X}_A(t) \perp\!\!\!\perp \mathcal{X}_B(t) \mid \mathcal{X}_S(t) \quad \forall t \in \mathbb{Z}.$$

Letting t tend to infinity, we can translate p -separation in the graph into conditional independence statements for complete subprocesses. For this, we define $\mathcal{X}_S(\infty) = \vee_{t \in \mathbb{Z}} \mathcal{X}_S(t)$ as the σ -algebra generated by the subprocess X_S .

Theorem 4.3. *Suppose X_V satisfies the block-recursive Granger-causal Markov property with respect to the graph G . Then, for any disjoint subsets A , B , and S of V , we have*

$$A \bowtie_p B \mid S \Rightarrow \mathcal{X}_A(\infty) \perp\!\!\!\perp \mathcal{X}_B(\infty) \mid \mathcal{X}_S(\infty).$$

We say that X satisfies the global AMP Markov property (GA) with respect to G .

Example 4.4. For an illustration of the global AMP Markov property, we consider again the graph G in Figure 2.1. In this graph, vertices 1 and 4 are not adjacent. Nevertheless, it can be shown that the two vertices cannot be p -separated by any set $S \subseteq \{2, 3, 5\}$: firstly, the path $1 \leftarrow 3 \rightarrow 4$ is p -connecting given a set S unless the set S contains the vertex 3 (Fig. 4.1 a). Secondly, the path $1 \rightarrow 3 \leftarrow 2 \leftarrow 4$ is p -connecting given S whenever both intermediate vertices 2 and 3 belong to S (Fig. 4.1 b). Finally, the path $1 \rightarrow 3 \leftarrow 2 \leftarrow 4$ is p -connecting given S if S contains vertex 3 but not 2 (Fig. 4.1 c). Thus, if X_V is a stationary process that obeys the block-recursive Granger-causal Markov property with respect to G , then the graph G does not encode that X_1 and X_4 are conditionally independent given X_S regardless of the choice of $S \subseteq \{2, 3, 5\}$.

Similarly, it can be shown that vertices 1 and 5 are p -separated given $S = \{3, 4\}$: every path between 1 and 5 that contains the edge $3 \rightarrow 5$ or the subpath $3 \rightarrow 4 \leftarrow 5$ is p -blocked by vertex 3. All other paths between 1 and 5 contain the subpath $2 \leftarrow 4 \leftarrow 5$ and, thus, are blocked by vertex 4. It follows that for every process

X_V that satisfies the block-recursive Granger-causal Markov property with respect to G the components X_1 and X_5 are conditionally independent given $X_{\{3,4\}}$.

4.2. The global Granger-causal Markov property

In this section, we apply the concept of pathwise separation to the problem of deriving general Granger noncausality relations from mixed graphs. To motivate the approach, we firstly consider the graphical VAR(1) model of all trivariate stationary processes $X_V = (X_1, X_2, X_3)$ given by

$$\begin{aligned} X_1(t+1) &= \phi_{11} X_1(t) + \phi_{12} X_2(t) + \varepsilon_1(t+1), \\ X_2(t+1) &= \phi_{22} X_2(t) + \phi_{23} X_3(t) + \varepsilon_2(t+1), \\ X_3(t+1) &= \phi_{33} X_3(t) + \varepsilon_3(t+1) \end{aligned} \quad (4.1)$$

for $t \in \mathbb{Z}$ with independent and standard normally distributed errors $\varepsilon_V(t)$, $t \in \mathbb{Z}$. The associated graph G that encodes the restrictions imposed on the model is shown in Figure 4.2. In this graph, the path $3 \rightarrow 2 \rightarrow 1$ is p -connecting given the empty set, which indicates that the components X_1 and X_3 are, in general, not independent in a bivariate analysis. However, an intuitive interpretation of the directed path $3 \rightarrow 2 \rightarrow 1$ suggests that X_3 Granger-causes X_1 but not vice versa if only the bivariate process $X_{\{1,3\}}$ is considered. Indeed, the block-recursive Granger-causal Markov property implies that

$$X_3(t+1) \perp\!\!\!\perp \mathcal{X}_{\{1,2\}}(t) \mid \mathcal{X}_{\{3\}}(t),$$

from which it follows by decomposition (see Appendix A) that

$$X_3(t+1) \perp\!\!\!\perp \mathcal{X}_{\{1\}}(t) \mid \mathcal{X}_{\{3\}}(t),$$

that is, X_1 is Granger-noncausal for X_3 with respect to $\mathcal{X}_{\{1,3\}}$. Obviously, the p -separation criterion is too strong for establishing this Granger-noncausality relationship between X_3 and X_1 since it requires that all paths between the two vertices are p -blocked whereas it seems sufficient that only certain paths, namely those ending with an arrowhead at vertex 3, are p -blocked.

This suggests the following definitions. A path π between two vertices a and b in G is said to be b -pointing if it has an arrowhead at the endpoint b . More generally, a path π between two disjoint subsets A and B is said to be B -pointing if it is b -pointing for some $b \in B$.

For the derivation of contemporaneous conditional independencies, we also need to consider paths with arrowheads at both endpoints; such paths π will be called *bi-pointing*. Furthermore, let $\pi = \langle \pi_1, \dots, \pi_n \rangle$ be a composition of paths π_i that are undirected or bi-pointing. Then π is said to be an *extended bi-pointing* path. In particular, this implies that any undirected or bi-pointing path is also an extended bi-pointing path; similarly, the composition $\pi = \langle \pi_1, \pi_2 \rangle$ of two extended bi-pointing




FIGURE 4.2. Mixed graph associated with trivariate time series in (4.1).

paths π_i is again extended bi-pointing. Moreover, every extended bi-pointing path π is of the form $\pi = \langle u_1, \beta, u_2 \rangle$ for some paths u_1 , u_2 , and β of possibly length zero, where u_1 and u_2 are undirected paths and β is a bi-pointing path (hence the term ‘extended bi-pointing’). With these definitions, we define the following global Granger-causal Markov property, which gives a path-oriented criterion for deriving general Granger noncausality relations from a mixed graph.

Definition 4.5 (Global Granger-causal Markov property). Let X_V be a stationary process and let $G = (V, E)$ be a mixed graph. Then X_V satisfies the *global Granger-causal Markov property* (GC) with respect to G if, for all disjoint subsets A , B , and S of V , the following conditions hold:

- (i) if every B -pointing path in G between A and B is p -blocked given $S \cup B$ then

$$X_A \nrightarrow X_B \ [\mathcal{X}_{A \cup B \cup S}];$$

- (ii) if every extended bi-pointing path in G between A and B is p -blocked given $A \cup B \cup S$ then

$$X_A \approx X_B \ [\mathcal{X}_{A \cup B \cup S}].$$

From the definition, it is immediately clear by setting $S = V \setminus (A \cup B)$ that the global Granger-causal Markov property entails the block-recursive Granger-causal Markov property. The following theorem shows that in fact, under conditions (M) and (P), the two Granger-causal Markov properties are equivalent; thus, the global Granger-causal Markov property may be employed to discuss the dynamic relationships implied by a graphical time series model defined in terms of the block-recursive Granger-causal Markov property.

Theorem 4.6. *Let X_V be a stationary process and let $G = (V, E)$ be a mixed graph. Then X_V satisfies the block-recursive Granger-causal Markov property with respect to G if and only if X_V satisfies the global Granger-causal Markov property with respect to G .*

As a consequence of the global Granger-causal Markov property, we find that p -separation in the graph implies Granger noncausality in both directions and contemporaneous conditional independence of the variables.

Corollary 4.7. *Suppose that the process X_V satisfies the block-recursive Granger-causal Markov property with respect to a mixed graph G . For disjoint subsets A , B , and S of V , if A and B are p -separated given S , then*

$$X_A \nrightarrow X_B \ [\mathcal{X}_{A \cup B \cup S}], \quad X_B \nrightarrow X_A \ [\mathcal{X}_{A \cup B \cup S}], \quad \text{and} \quad X_A \approx X_B \ [\mathcal{X}_{A \cup B \cup S}].$$

The following corollary summarizes the relationships between the various Markov properties for graphical time series models.

Corollary 4.8. *The various Granger-causal Markov properties are related as follows:*

$$(GC) \Leftrightarrow (BC) \Rightarrow (LC) \Leftrightarrow (PC).$$

If additionally condition (2.1) holds, then the four Granger-causal Markov properties are all equivalent. Furthermore, we have $(BC) \Rightarrow (GA)$

Proof. The corollary summarizes Theorems 2.5, 4.3, and 4.6. □




FIGURE 4.3. Illustration of global Granger-causal Markov property: Three 4-pointing paths (solid lines) between 1 and 4 that are p -blocked by the set $\{3, 4\}$.




FIGURE 4.4. Illustration of global Granger-causal Markov property: Three extended bi-pointing paths (solid lines) between 1 and 4 that are p -blocked by the set $\{3, 4\}$.

Example 4.9. For an illustration, we again consider a stationary time series X_V satisfying the block-recursive Granger-causal Markov property with respect to the graph G in Figure 2.1. In Example 4.4, we have seen that vertices 1 and 4 are not p -separated given $S = \{3\}$, that is, X_1 and X_4 are in general not conditionally independent given X_3 . We now employ the global Granger-causal Markov property to examine the dynamic relationships between the components X_1 and X_4 further.

Firstly, we note that all 4-pointing paths between 1 and 4 are p -blocked given the set $S = \{3, 4\}$. Three instances of such paths are depicted in Figure 4.3: while the first two paths are p -blocked by vertex 3, the last path is p -blocked by vertex 4 as an intermediate node. Straightforward considerations show that these three paths represent the three types of possible 4-pointing path ending either with $3 \rightarrow 4$, with $3 \rightarrow 5 \rightarrow 4$, or with $2 \leftarrow 4 \leftarrow 5 \rightarrow 4$, respectively. It follows that X_1 does not Granger-cause X_4 with respect to $\mathcal{X}_{\{1,3,4\}}$.

Similarly, we can examine all extended bi-pointing paths between vertices 1 and 4 to show that X_1 and X_4 are contemporaneously conditionally independent with respect to $\mathcal{X}_{\{1,3,4\}}$. Figure 4.4 shows three examples of such paths: the first two are p -blocked by vertex 3 (notice that on the second path, the vertex 3 is once a p -collider and once a p -noncollider) whereas the last path is p -blocked by vertices 2 and 4. For similar reasons as above, these three paths are exemplary for all extended bi-pointing paths between 1 and 4, and we conclude that X_1 and X_4 are indeed contemporaneously conditionally independent with respect to $\mathcal{X}_{\{1,3,4\}}$.

Finally, we note that every 1-pointing path between 4 and 1 must end with the directed edge $3 \rightarrow 1$. Since this edge has a tail at vertex 3, every such path must be p -blocked given $S = \{1, 3\}$, which implies that X_4 does not Granger-cause X_1 with respect to $\mathcal{X}_{\{1,3,4\}}$.

5. DISCUSSION

In this paper, we discussed a graphical modelling approach for multivariate time series that is based on mixed graphs in which each vertex represents one complete component series while the edges in the graph reflect possible dynamic interdependencies among the variables of the process. The constraints imposed by the graphs are formulated in terms of strong Granger noncausality and, thus, allow modelling

arbitrary non-linear dependencies. The graphical modelling approach can help to reduce the number of parameters involved in modelling high-dimensional non-linear time series while encoding the constraints on the parameters in a simple graph, which is easy to visualize and allows an intuitive understanding of the dependencies in the model.

We have shown that the interpretation of these graphs, which for many models are built only from pairwise Granger noncausality relations, is enhanced by so-called global Markov properties, which relate separation properties of the graph to conditional independence or Granger noncausality statements about the process. In this paper, we have used the path-oriented concept of p -separation, which allows to attribute Granger-causal relationships among the variables to certain pathways in the graphs.

Our objective has been to provide a general framework for modelling the dynamic interdependencies in multivariate time series; in particular, we focused on a simple graphical representation, which has been achieved by representing each component of a multivariate time series by a single vertex in the associated graph. The presented approach, however, is not the only possible, and since the first papers on the application of graphical models in time series analysis (Lynggaard and Walther 1993, Brillinger 1996), there has been an increasing interest in the topic (Stanghellini and Whittaker 1999, Dahlhaus 2000, Reale and Tunnicliffe Wilson 2001, Dahlhaus and Eichler 2003, Oxley et al. 2004, Moneta and Spirtes 2005, Eichler 2006a,b). All these approaches are basically restricted to the analysis of linear interdependencies, and most of them represent each variable at each time point by a separate vertex in the associated graph. In the following, we briefly compare our approach with alternative graphical representations and point out possible extensions.

Modelling processes of variables at separate time points

A more detailed modelling of dependencies among the components of a vector time series can be achieved by representing each random variable $X_v(t)$ by a different vertex v_t , say, in a graph G . This alternative approach has been discussed, for example, by Reale and Tunnicliffe Wilson (2001), Dahlhaus and Eichler (2003), and Moneta and Spirtes (2005). On the one hand, it leads to a more flexible class of graphical models and has the advantage that many of the concepts and methods that have been developed for the multivariate case carry over to the time series case. On the other hand, the increased flexibility leads to (sometimes much) larger graphs, which easily can become unwieldy and difficult to interpret, and it clearly also aggravates the model selection problem. Moreover, the underlying graph for such graphical time series models theoretically has infinitely many vertices, and it is not immediately clear how to prune this graph to a finite representation while preserving the Markov properties.

Apart from these theoretical and practical issues, we think that a high level of detail as provided by these models is not always wanted nor always appropriate. We give two examples. Firstly, Bacalá and Sameshima (2001) proposed a frequency-domain approach for the discussion of Granger-causal relationships based on the concept of partial directed coherence. Although this approach still requires the fitting of VAR models, the identification of interactions is performed in the frequency-domain

and hence only relations on the level of Granger noncausality can be identified. The results in Baccalá and Sameshima (2001) were summarized by path diagrams associated with the identified VAR model as discussed in Eichler (2006a). Our approach provides a theoretical framework for such analyses.

Secondly, multivariate time series are often obtained by high-frequency sampling of continuous-time processes such as EEG-recordings or neural spike trains. Here, our approach yields a graphical representation of the interrelationships that does not depend (to some extent) on the sampling frequency (e.g., Eichler 2005a). Moreover, many sophisticated models that have been proposed, for example, for analysing neural activity do not show a dependence on the past values only at specific lags. For instance, in the binary time series model discussed in Example 3.4, the conditional distribution of $X_b(t)$ given the past history $\mathcal{X}_V(t-1)$ depends on another process X_a through the past values $X_a(t-1), \dots, X_a(t-\gamma_b(t))$, where $\gamma_b(t)$ is the time elapsed since the last event of process X_b . In other words, the number of lagged variables $X_a(t-u)$ on which $X_b(t)$ depends varies over time depending on the past of X_b itself. Consequently, it seems inappropriate to break down the dependence of $X_a(t)$ on $\mathbf{X}_b(t)$ further into dependencies of $X_a(t)$ on $X_b(t-u)$ as required by the detailed modelling approach.

m-separation versus p-separation

The contemporaneous dependence structure of a process X_V can also be described by conditional independencies of the form

$$\mathcal{X}_A(t+1) \perp\!\!\!\perp \mathcal{X}_B(t+1) \mid \mathcal{X}_V(t),$$

in which case X_A and X_B are said to be contemporaneously independent with respect to $\mathcal{X}_V$. This alternative approach has been studied by (Eichler 2006a) in the context of weakly stationary processes and linear dependencies.

The most important difference between these two approaches for defining graphical time series models is that the corresponding composition and decomposition property

$$\begin{aligned} \mathcal{X}_A(t+1) \perp\!\!\!\perp \mathcal{X}_B(t+1) \mid \mathcal{X}_V(t) \\ \Leftrightarrow \mathcal{X}_a(t+1) \perp\!\!\!\perp \mathcal{X}_b(t+1) \mid \mathcal{X}_V(t) \quad \forall a \in A, \forall b \in B \end{aligned} \tag{5.1}$$

does not follow from conditions (M) and (P) but requires additional assumptions similarly to condition (2.1). Furthermore, we note that only the first two conditions in Proposition 3.1 are sufficient for the above property (5.1). Consequently, the class of graphical time series models for which the pairwise and the block-recursive Granger-causal Markov properties are equivalent would be smaller under the alternative approach based on contemporaneous independence.

Self-loops

In this paper, we have focused on modelling and analysing the interrelationships in multivariate time series. Therefore, we have not considered the possibility of directed self-loops $v \rightarrow v$, which could be used to impose additional constraints of the form $X_B(t+1) \perp\!\!\!\perp \mathcal{X}_B(t) \mid \mathcal{X}_{V \setminus B}(t)$ on a model. We note that, for a discussion of the dynamic interrelationships among variables, these self-loops are irrelevant. In fact, it can be shown that two disjoint sets A and B are p -separated given S in a

graph with self-loops if and only if they are also p -separated given S in the same graph with all self-loops removed. Similar statements can be formulated for pointing and extended bi-pointing paths.

Non-stationary time series

One of our main assumptions has been that the considered multivariate time series are stationary. This assumption, however, has been made mainly for the sake of simplicity, and the presented graphical modelling approach can be extended easily also to the case of non-stationary time series by requiring that the Granger noncausality and contemporaneous conditional independence constraints encoded by a graph hold at all time points in an interval $T \subseteq \mathbb{Z}$, say; in that case, we say that the time series obeys a Granger-causal Markov property with respect to the graph over the time interval T . This would allow to consider non-stationary time series models in which the pattern of dependencies remains fixed whereas the strength of the dependencies may change over time. An interesting extension would be models where also the graphical structure changes at certain times. For instance, Talih and Hengartner (2005) consider covariance selection models for multivariate time series where changes in the dependence structure occur at random times; this approach, however, does not model dynamic dependencies among the variables. Finally, we note that, from a statistical point of view, non-stationary models are of much less interest than stationary models due to the involved practical problems.

Two important issues have not been addressed in this paper. Firstly, in many applications there is little prior knowledge about the causal relationships between the variables, and empirical methods have to be used to find an appropriate graphical model. This step of model selection is hampered by the large number of possible models by which an exhaustive search becomes infeasible even for moderate dimensions. Therefore, model search strategies are required to lessen the computational burden.

A second issue, which is related to the problem of model selection, is the identification of causal effects. It is clear from the definition of Granger causality that we may conclude from Granger causality to the existence of a causal effect only if all relevant variables are included in a study, whereas the omission of important variables can lead to spurious causalities. However, Hsiao (1982) noted that such spurious causalities may vanish if the information set is reduced. In other words, two processes that both satisfy the pairwise causal Markov property with respect to a graph G may exhibit different Granger noncausality relations with respect to partial information sets due to the presence or absence of spurious causalities. Some concepts how this observation could be exploited for causal inference have been discussed in Eichler (2005a,b).

APPENDIX A. CONDITIONAL INDEPENDENCE AND STOCHASTIC PROCESSES

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\mathcal{F}_1$, $\mathcal{F}_2$, and $\mathcal{F}_3$ sub- σ -algebras of $\mathcal{F}$. Here and elsewhere we assume for simplicity that all sub- σ -algebras are completed¹. The smallest σ -algebra generated by $\mathcal{F}_i \cup \mathcal{F}_j$ is denoted as $\mathcal{F}_i \vee \mathcal{F}_j$. Then $\mathcal{F}_1$ and $\mathcal{F}_2$

¹Otherwise the σ -algebras on both sides of (A.1) need to be completed explicitly.

are said to be independent conditionally on $\mathcal{F}_3$ if $\mathbb{E}(X|\mathcal{F}_2 \vee \mathcal{F}_3) = \mathbb{E}(X|\mathcal{F}_3)$ a.s. for all real-valued, bounded, $\mathcal{F}_1$ -measurable random variables X . Using the notation of Dawid (1979) we write $\mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_2 | \mathcal{F}_3$ $[\mathbb{P}]$ or $\mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_2 | \mathcal{F}_3$ if the reference to $\mathbb{P}$ is clear.

Let $\mathcal{F}_i$, $i = 1, \dots, 4$ be sub- σ -algebras of $\mathcal{F}$. Then the basic properties of the conditional independence relation are:

- (CI1) $\mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_2 | \mathcal{F}_3 \Rightarrow \mathcal{F}_2 \perp\!\!\!\perp \mathcal{F}_1 | \mathcal{F}_3$ (symmetry)
- (CI2) $\mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_2 \vee \mathcal{F}_3 | \mathcal{F}_4 \Rightarrow \mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_2 | \mathcal{F}_4$ (decomposition)
- (CI3) $\mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_2 \vee \mathcal{F}_3 | \mathcal{F}_4 \Rightarrow \mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_2 \vee \mathcal{F}_3 | \mathcal{F}_3 \vee \mathcal{F}_4$ (weak union)
- (CI4) $\mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_2 | \mathcal{F}_4$ and $\mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_3 | \mathcal{F}_2 \vee \mathcal{F}_4 \Rightarrow \mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_2 \vee \mathcal{F}_3 | \mathcal{F}_4$ (contraction)

In some of the proofs in this paper, we make use of an additional property,

- (CI5) $\mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_2 | \mathcal{F}_3 \vee \mathcal{F}_4$ and $\mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_3 | \mathcal{F}_2 \vee \mathcal{F}_4 \Leftrightarrow \mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_2 \vee \mathcal{F}_3 | \mathcal{F}_4$,

which has been called *intersection property* by Pearl (1988). Unlike the other basic properties of conditional indepenence, this property does not hold in general. A sufficient and necessary condition for (CI5) is given by

$$(\mathcal{F}_2 \vee \mathcal{F}_4) \cap (\mathcal{F}_3 \vee \mathcal{F}_4) = \mathcal{F}_4. \quad (\text{A.1})$$

In that case, $\mathcal{F}_2$ and $\mathcal{F}_3$ are said to be measurably separated conditionally on $\mathcal{F}_4$, denoted by $\mathcal{F}_2 \parallel \mathcal{F}_3 | \mathcal{F}_4$ (Florens et al. 1990). We note that conditional measurable separability preserves most, but not all, of the properties of conditional independence. For instance, if for some sub- σ -algebras $\mathcal{F}_i$, $i = 1, \dots, 4$ of $\mathcal{F}$ we have $\mathcal{F}_1 \parallel \mathcal{F}_2 | \mathcal{F}_3$ and $\mathcal{F}_4 \subseteq \mathcal{F}_2$ then $\mathcal{F}_1 \parallel \mathcal{F}_4 | \mathcal{F}_3$. For details, we refer to Chapter 5.2 of Florens et al. (1990).

If $\mathcal{F}_i$ is generated by random vectors X_i for $i = 1, \dots, 4$, then a sufficient condition for conditional measurable separability of the X_i 's and, thus, of the $\mathcal{F}_i$'s is that the probability measure $\mathbb{P}^{X_1, \dots, X_4}$ is absolutely continuous with respect to a product measure μ and has a positive and continuous density. However, if each of the σ -algebras $\mathcal{F}_i$ is generated by infinitely many random variables, the condition is obviously no longer valid. Therefore, we show that, in the case of stationary processes X_V , the above assumption of existence of a positive and continuous density need only be made for the conditional distribution of $X_V(t+1)$ given its past $\mathbf{X}_V(t)$ if the process is strongly mixing.

Lemma A.1. *Suppose that X_V is a stationary process such that condition (P) holds, and let Y_1, Y_2 be finite disjoint subsets of $S(t) = \{X_v(s), s \leq t, v \in V\}$. Then*

$$Y_1 \parallel Y_2 | \sigma(S(t) \setminus (Y_1 \cup Y_2)). \quad (\text{A.2})$$

Proof. A sufficient condition for (A.2) (Florens et al. 1990, Corollary 5.2.11) is the existence of a probability measure $\mathbb{P}'$ on $(\Omega, \mathcal{X}_V(t))$ such that $\mathbb{P}|_{\mathcal{X}_V(t)}$ and $\mathbb{P}'$ are equivalent (i.e. have the same null sets) and

$$Y_1 \perp\!\!\!\perp Y_2 | \sigma(S(t) \setminus (Y_1 \cup Y_2)) \quad [\mathbb{P}']. \quad (\text{A.3})$$

Let $Z_j = X_V(t-j)$ for $j = 0, \dots, k-1$ and $Z_k = S(t-k)$. Noting that the conditional densities $f_{Z_{jv}|Z_k}$ exist and can be derived from the product of the conditional

densities $f_{Z_j|Z_{j+1},\dots,Z_k}$, we define the probability kernel $Q(z_k, A)$ from $\mathbb{R}^{V \times \mathbb{N}}$ to $\mathbb{R}^{V \times k}$ by

$$Q(z_k, A_0 \times \dots \times A_{k-1}) = \int_{A_0} \dots \int_{A_{k-1}} \prod_{j=0}^{k-1} \prod_{v \in V} f_{Z_{jv}|Z_k}(z_{jv}|z_k) \nu^k(dz_0, \dots, dz_{k-1}).$$

Then the probability $\mathbb{P}'$ on $\mathcal{X}_V(t)$ defined by

$$\mathbb{P}'(Z_j \in A_j, j = 0, \dots, k) = \int_{Z_k^{-1}(A_k)} Q(Z_k, (dz_0, \dots, dz_{k-1})) d\mathbb{P}$$

is equivalent to $\mathbb{P}$ and the random variables Z_{jv} for $j = 0, \dots, k-1$ and $v \in V$ are mutually conditionally independent given Z_k . This implies in particular (A.3) for k large enough. $\square$

The next result shows that for strongly mixing processes this conditional measurable separability can also be extended to σ -algebras $\mathcal{X}_A(t)$ generated by the pasts $\mathbf{X}_A(t)$.

Proposition A.2. *Suppose that X_V is a stationary process such that conditions (M) and (P) hold. Then $\mathcal{X}_A(t)$ and $\mathcal{X}_B(t)$ are measurably separated conditionally on $\mathcal{X}_{V \setminus (A \cup B)}(t)$ for all disjoint subsets A and B of V .*

For the proof of the proposition, we firstly show that every stationary strongly mixing process is also conditionally mixing. Let $A \subseteq V$. For any set $S' = \{\omega \in \Omega \mid \mathbf{X}_A(t-k-1) \in T\}$ in $\mathcal{X}_A(t-k-1)$, we define the shifted set $S'_j \in \mathcal{X}_A(t-k-j)$ by $S'_j = \{\omega \in \Omega \mid \mathbf{X}_A(t-k-j) \in T\}$. Then the process X_V is said to be conditionally mixing (Veijanen 1990) if for all disjoint subsets A and B of V , $t \in \mathbb{Z}$, $S \in \sigma\{X_A(t), \dots, X_A(t-k)\}$, and $S' \in \mathcal{X}_A(t-k-1)$ we have

$$\mathbb{E}|\mathbb{P}(S \cap S'_j \mid \mathcal{X}_B(t)) - \mathbb{P}(S \mid \mathcal{X}_B(t))\mathbb{P}(S'_j \mid \mathcal{X}_B(t))| \rightarrow 0$$

as $j \rightarrow \infty$.

Lemma A.3. *Let X_V be a stationary process such that condition (M) holds. Then X_V is also conditionally mixing.*

Proof. Let $F \in \sigma\{X_B(t_1), \dots, X_B(t_m)\}$ with $m \in \mathbb{N}$ and $t_j < t$, $1 \leq j \leq m$. Noting that

$$\mathbb{E}[\mathbb{E}(1_S \mid \mathcal{X}_B(t))\mathbb{E}(1_{S'_j} \mid \mathcal{X}_B(t)) \cdot 1_F] = \mathbb{E}[\mathbb{E}(1_{S \cap F} \mid \mathcal{X}_B(t)) \cdot 1_{S'_j}]$$

by the law of iterated expectation by conditioning on $\mathcal{X}_B(t)$, we have for $S \in \sigma\{X_A(t), \dots, X_A(t-k)\}$ and $S' \in \mathcal{X}_A(t-k-1)$,

$$\begin{aligned} & |\mathbb{E}[\{\mathbb{E}(1_{S \cap S'_j} \mid \mathcal{X}_B(t)) - \mathbb{E}(1_S \mid \mathcal{X}_B(t))\mathbb{E}(1_{S'_j} \mid \mathcal{X}_B(t))\} \cdot 1_F]| \\ & \leq |\mathbb{E}(1_{S \cap S'_j \cap F}) - \mathbb{E}(1_{S \cap F})\mathbb{E}(1_{S'_j})| \\ & \quad + |\mathbb{E}[\mathbb{E}(1_{S \cap F} \mid \mathcal{X}_B^{(n)}(t))]\mathbb{E}(1_{S'_j}) - \mathbb{E}[\mathbb{E}(1_{S \cap F} \mid \mathcal{X}_B^{(n)}(t)) \cdot 1_{S'_j}]| \\ & \quad + |\mathbb{E}[\{\mathbb{E}(1_{S \cap F} \mid \mathcal{X}_B^{(n)}(t)) - \mathbb{E}(1_{S \cap F} \mid \mathcal{X}_B(t))\} \cdot 1_{S'_j}]| \end{aligned}$$

where $\mathcal{X}_B^{(n)}(t) = \sigma\{X_B(t), \dots, X_B(t-n)\}$. Each of the three terms on the right side converges to zero as $j-n$ and n tend to infinity. Since the union of all $\sigma\{X_B(t_1), \dots, X_B(t_m)\}$ with $m \in \mathbb{N}$ and $t_j < t$ constitutes a $\cap$ -stable generator of $\mathcal{X}_B(t)$, the assertion of the lemma follows. $\square$

Proof of Proposition A.2. Let A and B be disjoint subsets of V . We have to show that $\mathcal{X}_A(t)$, $\mathcal{X}_B(t)$, and $\mathcal{X}_{V \setminus (A \cup B)}(t)$ satisfy (A.1) and hence that

$$\mathcal{X}_{V \setminus B}(t) \cap \mathcal{X}_{V \setminus A}(t) = \mathcal{X}_{V \setminus (A \cup B)}(t) \quad (\text{A.4})$$

for all $t \in \mathbb{Z}$. From Lemma A.1, it follows that, for all $t \in \mathbb{Z}$ and $k \in \mathbb{N}$, the σ -algebras $\sigma\{X_A(t), \dots, X_A(t-k+1)\}$ and $\sigma\{X_B(t), \dots, X_B(t-k+1)\}$ are measurably separable conditional on $\mathcal{X}_{V \setminus (A \cup B)}(t) \vee \mathcal{X}_V(t-k)$. Accordingly, we have by the definition of conditionally measurable separability

$$(\mathcal{X}_{V \setminus B}(t) \vee \mathcal{X}_V(t-k)) \cap (\mathcal{X}_{V \setminus A}(t) \vee \mathcal{X}_V(t-k)) = \mathcal{X}_{V \setminus (A \cup B)}(t) \vee \mathcal{X}_V(t-k)$$

for all $t \in \mathbb{Z}$ and $k \in \mathbb{N}$. Since the σ -algebras on both sides are monotonically decreasing as k increases, this yields for $k \rightarrow \infty$

$$\begin{aligned} \bigcap_{k>0} [(\mathcal{X}_{V \setminus B}(t) \vee \mathcal{X}_V(t-k)) \cap (\mathcal{X}_{V \setminus A}(t) \vee \mathcal{X}_V(t-k))] \\ = \bigcap_{k>0} (\mathcal{X}_{V \setminus (A \cup B)}(t) \vee \mathcal{X}_V(t-k)) \end{aligned}$$

for all $t \in \mathbb{Z}$. In order to establish (A.4), it therefore suffices to show that

$$\bigcap_{k>0} [\mathcal{X}_S(t) \vee \mathcal{X}_V(t-k)] = \mathcal{X}_S(t) \quad (\text{A.5})$$

for any subset S of V . Let $\mathcal{T}$ denote the tail σ -algebra on the left hand side. Since by Lemma A.3 the process X_V is conditionally mixing, the σ -algebras $\mathcal{T}$ and $\mathcal{X}_{V \setminus S}^{(k)}(t) = \sigma\{X_{V \setminus S}(t), \dots, X_{V \setminus S}(t-k)\}$ are conditionally independent given $\mathcal{X}_S(t)$. Thus, we have for all $T \in \mathcal{T}$

$$\mathbb{E}(1_T | \mathcal{X}_S(t)) = \mathbb{E}(1_T | \mathcal{X}_{V \setminus S}^{(k)}(t) \vee \mathcal{X}_S(t)) \rightarrow 1_T$$

as $k \rightarrow \infty$. Hence $\mathcal{T} \subseteq \mathcal{X}_S(t)$ whereas the converse relation is obvious. $\square$

Proof of Proposition 2.4. The result follows from Proposition A.2 and the fact that (A.1) is a necessary and sufficient condition for (CI5). $\square$

APPENDIX B. GRAPHICAL TERMINOLOGY

We firstly recall some basic graphical definitions used in this paper. In a graph $G = (V, E)$, if there is a directed edge $a \rightarrow b$, we say that a is a parent of b and b is a child of a ; similarly, if there is an undirected line $a - b$, the vertices a and b are called neighbours. The sets of parents, children and neighbours of a vertex a are denoted as $\text{pa}(a)$, $\text{ch}(a)$, and $\text{ne}(a)$, respectively. Furthermore, for $A \subseteq V$, let $\text{pa}(A) = \cup_{a \in A} \text{pa}(a) \setminus A$ be the set of all parents of vertices in A that are not themselves in A , and let $\text{ch}(A)$ and $\text{ne}(A)$ be defined similarly.

Next, as in Frydenberg (1990), a node b is said to be an *ancestor* of a if either $b = a$ or there exists a directed path $b \rightarrow \dots \rightarrow a$ in G . The set of all ancestors of elements in A is denoted by $\text{an}(A)$. Notice that this definition differs from the one given in Lauritzen (1996). A subset A is called an *ancestral set* if it contains all its ancestors, that is, $\text{an}(A) = A$.

Finally, let $G = (V, E)$ and $G' = (V', E')$ be mixed graphs. Then G' is a *subgraph* of G if $V' \subseteq V$ and $E' \subseteq E$. If A is a subset of V it induces the subgraph $G_A = (A, E_A)$ where E_A contains all edges $e \in E$ that have both endpoints in A .

In the remainder of this section, we prove some auxiliary results that allow us to relate separation statements in the full graph to separation statement in so-called marginal ancestral graphs, which basically reflect the dynamic dependencies in appropriate subprocesses (see Lemma C.1).

Definition B.1 (Marginal ancestral graph). Let $G = (V, E)$ be a mixed graph and let A be an ancestral subset of V . Then the marginal ancestral graph $G_{[A]} = (A, E_{[A]})$ induced by A is obtained from the induced subgraph G_A by insertion of additional undirected edges $a - b$ whenever there exists an undirected path between a and b in G that does not intersect $\text{an}(A) \setminus \{a, b\}$.

Lemma B.2. *Let $G = (V, E)$ be a mixed graph and A, B, S disjoint subsets of V . Then A and B are p -separated given S in G if and only if A and B are p -separated given S in $G_{[\text{an}(A \cup B \cup S)]}$.*

Proof. To show necessity, let $\pi = \langle e_1, \dots, e_n \rangle$ be a p -connecting path between A and B given S in $G_{[\text{an}(A \cup B \cup S)]}$. If all edges of π are edges in G , π is also p -connecting given S in G . Thus, we may assume that the edges $e_{j_1}, \dots, e_{j_m}$ in π do not occur in G . These edges e_{j_k} are necessarily undirected since all directed edges in $G_{[\text{an}(A \cup B \cup S)]}$ also occur in G . Let $e_{j_k} = v_{j_k} - v_{j_k+1}$. Then by definition of the marginal ancestral graph there exists an undirected path ϕ_{j_k} between v_{j_k-1} and v_{j_k} which bypasses $\text{an}(A \cup B \cup S) \setminus \{v_{j_k-1}, v_{j_k}\}$ and therefore is p -connecting given S . Replacing all edges e_{j_k} in π by the corresponding paths ϕ_{j_k} we obtain a new path π' which connects A and B in G . This path π' is also p -connecting given S since the replacement of e_{j_k} by the undirected and p -connecting path ϕ_{j_k} does not change the p -collider resp. p -noncollider status of the nodes v_{j_k-1} and v_{j_k} .

Conversely for sufficiency, let $\pi = \langle e_1, \dots, e_n \rangle$ be a p -connecting path between A and B given S in G . Then all edges in π with both endpoints in $\text{an}(A \cup B \cup S)$ also occur in $G_{[\text{an}(A \cup B \cup S)]}$ since $G_{\text{an}(A \cup B \cup S)}$ is a subgraph of $G_{[\text{an}(A \cup B \cup S)]}$. We firstly show that the endpoints of any directed edge e_j in π are in $\text{an}(A \cup B \cup S)$. Let $e_j = v_j \rightarrow v_{j+1}$ (the case $e_j = v_j \leftarrow v_{j+1}$ is treated similarly). Then there exists a directed subpath $\langle e_j, \dots, e_{j+r} \rangle$ of maximal length such that either v_{j+r} is an endpoint of π and, thus, in $A \cup B$ or e_{j+r+1} is of the form $v_{j+r} - v_{j+r+1}$ or $v_{j+r} \leftarrow v_{j+r+1}$. In the latter case v_{j+r} is a p -collider and, thus, in S since π is p -connecting given S . It follows that v_j and v_{j+1} are both in $\text{an}(A \cup B \cup S)$.

Next, if e_j is an edge in π that does not occur in $G_{[\text{an}(A \cup B \cup S)]}$, at least one of its endpoints v_{j-1} and v_j is not in $\text{an}(A \cup B \cup S)$. Thus, there exists an undirected subpath $\psi_{i,k} = \langle e_i, \dots, e_k \rangle$ with $i \leq j \leq k$ such that $v_{i-1}, v_k \in \text{an}(A \cup B \cup S)$ but all intermediate points are not in $\text{an}(A \cup B \cup S)$. In other words, v_{i-1} and v_k are not separated by $\text{an}(A \cup B \cup S) \setminus \{v_{j-1}, v_k\}$ in G which implies the presence of the undirected edge $f_{i,k} = v_{i-1} - v_k$ in $G_{[\text{an}(A \cup B \cup S)]}$. Replacing all undirected subpaths $\phi_{i,k}$ with intermediate points not in $\text{an}(A \cup B \cup S)$ by the corresponding edge $f_{i,k}$, we obtain a path between A and B in $G_{[\text{an}(A \cup B \cup S)]}$ which still has all its p -collider in S and all its p -noncolliders outside S and therefore is p -connecting given S . $\square$

The following lemma is an adapted version of Proposition 2 in Koster (1999). The proof is considerably shorter due to the fact that we allow paths to be self-intersecting.

Lemma B.3. *Let A, B, S be disjoint subsets of V . Then A and B are p -separated given S in $G_{[\text{an}(A \cup B \cup S)]}$ if and only there exist subsets A' and B' such that $A \subseteq A'$, $B \subseteq B'$, $A' \cup B' \cup S = \text{an}(A \cup B \cup S)$ and*

$$A' \not\propto_p B' \mid S \text{ } [G_{[\text{an}(A \cup B \cup S)]}].$$

Proof. By Lemma B.2 we may assume that $V = \text{an}(A \cup B \cup S)$. Let A' be the subset of vertices $v \in V \setminus (B \cup S)$ such that $v \not\propto_p B \mid S \text{ } [G]$, and set $B' = V \setminus (A' \cup S)$. Then A' and B are obviously p -separated given S . Thus, we have to show that a and b' are p -separated given S whenever $a \in A'$ and $b' \in B' \setminus B$. Suppose to the contrary that there exists a p -connecting path π between some $a \in A'$ and $b' \in B' \setminus B$. Since A' contains all vertices in $V \setminus (B \cup S)$ that are p -separated from B given S , there exists a p -connecting path π' between b' and some $b \in B$. Furthermore, since $b' \in \text{an}(A \cup B \cup S) \setminus (A \cup B \cup S)$ there exists some vertex $u \in A \cup B \cup S$ and a directed path $\omega = b \rightarrow \dots \rightarrow u$ with no intermediate vertices in $A \cup B \cup S$. Denoting by $\bar{\omega}$ the reverse path of ω , that is, $\bar{\omega} = u \leftarrow \dots \leftarrow b$, we may compose a path ϕ between A and B by

- (i) $\phi = \langle \bar{\omega}, \pi' \rangle$ if $v \in A$,
- (ii) $\phi = \langle \pi, \omega \rangle$ if $v \in B$, and
- (iii) $\phi = \langle \pi, \omega, \bar{\omega}, \pi' \rangle$ if $v \in S$.

We note that the directed path ω is p -connecting given S since it has no intermediate vertices in S . Furthermore, $b' \notin S$ is a p -noncollider on ϕ in each of these cases and $v \in S$ is a p -collider on ϕ in case (iii). Hence ϕ is a p -connecting path between A and B given S which contradicts our assumption.

The opposite implication is obvious because of the elementwise definition of p -separation. $\square$

Because of Lemmas B.2 and B.3, it is often sufficient in the proofs to consider only the case of $A \not\propto_p B \mid S$ with $S = V \setminus (A \cup B)$. In this case, p -separation can be characterized in terms of pure-collider paths—paths on which every intermediate node is a collider—or in terms of local configurations.

Lemma B.4. *Let G be a mixed graph and let A and B be two disjoint subsets of V . Then the following statements are equivalent:*

- (i) $A \not\propto_p B \mid V \setminus (A \cup B)$;
- (ii) A and B are not connected by a pure-collider path;
- (iii) $(A \cup \text{ch}(A)) \cap (B \cup \text{ch}(B)) = \emptyset$ and $\text{ne}(A \cup \text{ch}(A)) \cap (B \cup \text{ch}(B)) = \emptyset$.

Proof. This observation follows directly from the definition of p -separation and pure-collider paths. $\square$

APPENDIX C. PROOFS

Proof of Theorem 2.5. Setting $A = \{a\}$ in (BC), we immediately obtain (LC). Conversely, since $\text{pa}(a) \cup \{a\} \subseteq \text{pa}(A) \cup A$, we have by (LC) together with (CI2) and (CI3)

$$X_{V \setminus \text{pa}(A) \cup A} \not\leftrightarrow X_a \quad \forall a \in A,$$

which, under condition (2.1), implies the first part of (BC). The second part is proved similarly.

To see that (LC) and (PC) are equivalent, we note that, under conditons (M) and (P), the intersection property leads to the following composition and decomposition property for Granger noncausality relations:

$$X_A \nrightarrow X_B [\mathcal{X}_V] \Leftrightarrow X_a \nrightarrow X_B [\mathcal{X}_V] \quad \forall a \in A. \quad (\text{C.1})$$

Similarly, we have for contemporaneous conditional independence relations

$$X_A \approx X_B [\mathcal{X}_V] \Leftrightarrow X_a \approx X_b [\mathcal{X}_V] \quad \forall a \in A, \forall b \in B. \quad (\text{C.2})$$

Taking $A = V \setminus (B \cup \text{pa}(B))$ in (C.1) and $A = V \setminus (B \cup \text{pa}(B))$ in (C.2), we find that the pairwise and the local Granger-causal Markov property are equivalent. $\square$

Proof of Proposition 3.1. We have to show that each of the three conditions (i), (ii), and (iii) implies

$$X_A \nrightarrow X_b [\mathcal{X}_V] \quad \forall b \in B \Rightarrow X_A \nrightarrow X_B [\mathcal{X}_V] \quad (\text{C.3})$$

for any two disjoint subsets $A, B \subseteq V$.

For the first case, let H be the Hilbert space of all square integrable random variables on $(\Omega, \mathcal{F}, \mathbb{P})$. Furthermore, for $U \subseteq V$, let $H_U(t)$ be the closed subspace spanned by $\{X_u(s), u \in U, s \leq t\}$ and let $H_U^\perp(t)$ be its orthogonal complement. Then we have for any $Y \in H_{V \setminus A}^\perp(t)$

$$\text{cov}(X_B(t+1), Y) = 0 \Leftrightarrow \text{cov}(X_b(t+1), Y) = 0 \quad \forall b \in B,$$

which for a Gaussian process implies (C.3).

Next, suppose that condition (ii) holds and that X_A is Granger-noncausal for X_b with respect to $\mathcal{X}_V$ for all $b \in B$. Then, the conditional distribution $\mathbb{P}^{X_B(t+1)|\mathbf{X}_V(t)}$ satisfies

$$\mathbb{P}^{X_B(t+1)|\mathbf{X}_V(t)} = \otimes_{b \in B} \mathbb{P}^{X_b(t+1)|\mathbf{X}_V(t)} = \otimes_{b \in B} \mathbb{P}^{X_b(t+1)|\mathbf{X}_{V \setminus A}(t)}$$

and, thus, is $\mathcal{X}_{V \setminus A}(t)$ -measurable, which proves (C.3).


Finally, if condition (iii) holds, we have

$$X_B(t+1) - \mathbb{E}[X_B(t+1) | \mathcal{X}_V(t)] \perp\!\!\!\perp \mathcal{X}_A(t) | \mathcal{X}_{V \setminus A}(t).$$

Since the left hand side of (C.3) implies that $\mathbb{E}[X_B(t+1) | \mathcal{X}_V(t)]$ is $\mathcal{X}_{V \setminus A}(t)$ -measurable, we obtain $X_B(t+1) \perp\!\!\!\perp \mathcal{X}_A(t) | \mathcal{X}_{V \setminus A}(t)$, which completes the proof. $\square$

For the proofs of the equivalence of the block-recursive and the global Granger-causal Markov property, it will be convenient to restrict ourselves to mixed graphs for ancestral subsets. Due to the additional undirected edges inserted into the marginal ancestral graph $G_{[\text{an}(A)]}$, the subprocess $X_{\text{an}(A)}$ satisfies the pairwise Granger-causal Markov property with respect to $G_{[\text{an}(A)]}$ if X_V did so with respect to G . The following lemma shows that the same inheritance property also holds for the block-recursive Granger-causal Markov property.

Lemma C.1. *Suppose that X_V satisfies the block-recursive Granger-causal Markov property with respect to the mixed graph G , and let $U \subseteq V$. Then the subprocess $X_{\text{an}(U)}$ satisfies the block-recursive Granger-causal Markov property with respect to the marginal ancestral graph $G_{[\text{an}(U)]}$.*


 FIGURE C.1. Pure-collider paths between two vertices a and b .

Proof. Let $H = G_{[\text{an}(U)]}$ and let A be a subset of $\text{an}(U)$. We firstly note that, since $\text{an}(U)$ is an ancestral set and, thus, contains the parents of all its subsets A , the parents of A in both graphs are the same, i.e. $P = \text{pa}_G(A) = \text{pa}_H(A)$. By the block-recursive Granger-causal Markov property of X_V with respect to G , $X_{V \setminus (P \cup A)}$ does not Granger-cause X_A with respect to $\mathcal{X}_V$, which by (CI2) implies that $X_{\text{an}(U) \setminus (P \cup A)}$ is Granger-noncausal for X_A with respect to the smaller filtration $\mathcal{X}_{\text{an}(U)}$ as required by the block-recursive Granger-causal Markov property of $X_{\text{an}(U)}$ with respect to H .

Next, let $N = \text{ne}_H(A)$. Then A and $\text{an}(U) \setminus N$ are separated by N in H^u , that is, a and b are not adjacent in the undirected subgraph H^u whenever $a \in A$ and $b \in \text{an}(U) \setminus N$. By definition of H , this implies that A and $\text{an}(U) \setminus N$ are separated by N in G^u . By the block-recursive Granger-causal Markov property, it follows that

$$\mathcal{X}_A(t+1) \perp\!\!\!\perp \mathcal{X}_{\text{an}(U) \setminus N}(t+1) \mid \mathcal{X}_N(t+1) \vee \mathcal{X}_V(t)$$

and

$$\mathcal{X}_A(t+1) \perp\!\!\!\perp \mathcal{X}_{V \setminus \text{an}(U)}(t) \mid \mathcal{X}_{\text{an}(U)}(t).$$

Combining these two relations, we find that X_A and $X_{\text{an}(U) \setminus N}$ are contemporaneously conditionally independent with respect to $\mathcal{X}_{\text{an}(U)}$ as required by the block-recursive Granger-causal Markov property of $X_{\text{an}(U)}$ with respect to the graph H . $\square$

Proof of Lemma 4.2. For notational convenience, we may assume in view of Lemma C.1 that $\text{an}(A \cup B \cup S) = V$ and, thus, $G_{[\text{an}(A \cup B \cup S)]} = G$. Furthermore, Lemma B.2 implies that, if $A \bowtie_p B \mid S$ in the graph G , there exists a partition (A^*, B^*, S) of V such that $A \subseteq A^*$, $B \subseteq B^*$, and $A^* \bowtie_p B^* \mid S$. Thus, without loss of generality, we may assume that $S = V \setminus (A \cup B)$.

With these simplifications, it suffices to show that $A \bowtie_p B \mid V \setminus (A \cup B)$ implies

$$\mathcal{X}_{X_A}(t) \perp\!\!\!\perp \mathcal{X}_{X_B}(t) \mid \mathcal{X}_{V \setminus (A \cup B)}(t) \quad (\text{C.4})$$

for all $t \in \mathbb{Z}$. To this end, we firstly show that

$$\mathcal{X}_A(t) \perp\!\!\!\perp \mathcal{X}_B(t) \mid \mathcal{X}_{V \setminus (A \cup B)}(t) \vee \mathcal{X}_{A \cup B}(t-k) \quad (\text{C.5})$$

for all $t \in \mathbb{Z}$ and $k \in \mathbb{N}$.

We proceed by induction on k . For $k = 1$, we note that $B \subseteq V \setminus (A \cup \text{ne}(A))$ and hence

$$\mathcal{X}_A(t) \perp\!\!\!\perp \mathcal{X}_B(t) \mid \mathcal{X}_{V \setminus (A \cup B)}(t) \vee \mathcal{X}_V(t-1). \quad (\text{C.6})$$

For the induction step $k \rightarrow k+1$ assume that

$$\mathcal{X}_A(t) \perp\!\!\!\perp \mathcal{X}_B(t) \mid \mathcal{X}_{V \setminus (A \cup B)}(t) \vee \mathcal{X}_{A \cup B}(t-k) \quad (\text{C.7})$$

for all $t \in \mathbb{Z}$. Let $C_A = A \cup \text{ch}(A)$. Then, since by the block-recursive Granger-causal Markov property X_A is Granger-noncausal for $X_{V \setminus C_A}$, we have

$$\mathcal{X}_A(t) \perp\!\!\!\perp \mathcal{X}_{V \setminus C_A}(t+1) \mid \mathcal{X}_{V \setminus A}(t) \vee \mathcal{X}_{A \cup B}(t-k)$$

and further with (C.7) and (CI4)

$$\mathcal{X}_A(t) \perp\!\!\!\perp \mathcal{X}_B(t) \vee \mathcal{X}_{V \setminus C_A}(t+1) \mid \mathcal{X}_{V \setminus (A \cup B)}(t) \vee \mathcal{X}_V(t-k).$$

With $N_A = \text{ne}(A \cup \text{ch}(A)) = \text{ne}(C_A)$, we obtain by (CI2) and (CI3)

$$\mathcal{X}_A(t) \perp\!\!\!\perp \mathcal{X}_B(t) \vee \mathcal{X}_{V \setminus (C_A \cup N_A)}(t+1) \mid \mathcal{X}_{N_A}(t+1) \vee \mathcal{X}_{V \setminus (A \cup B)}(t) \vee \mathcal{X}_V(t-k). \quad (\text{C.8})$$

Next, we note that by Lemma B.4 $B \cup \text{ch}(B) \subseteq V \setminus (C_A \cup N_A)$ and thus

$$\mathcal{X}_{C_A}(t+1) \perp\!\!\!\perp \mathcal{X}_B(t) \mid \mathcal{X}_{N_A}(t+1) \vee \mathcal{X}_{V \setminus B}(t).$$

Furthermore, X_{C_A} and $X_{V \setminus (C_A \cup N_A)}$ are contemporaneously conditionally independent and thus

$$\mathcal{X}_{C_A}(t+1) \perp\!\!\!\perp \mathcal{X}_{V \setminus (C_A \cup N_A)}(t+1) \mid \mathcal{X}_{N_A}(t+1) \vee \mathcal{X}_V(t).$$

Together with the previous relation, we obtain by (CI4)

$$\mathcal{X}_{C_A}(t+1) \perp\!\!\!\perp \mathcal{X}_B(t) \vee \mathcal{X}_{V \setminus (C_A \cup N_A)}(t+1) \mid \mathcal{X}_{N_A}(t+1) \vee \mathcal{X}_{V \setminus B}(t).$$

By (C.8) and the intersection property (CI5), this yields

$$\mathcal{X}_A \vee \mathcal{X}_{C_A}(t+1) \perp\!\!\!\perp \mathcal{X}_B(t) \vee \mathcal{X}_{V \setminus (C_A \cup N_A)}(t+1) \mid \mathcal{X}_{N_A}(t+1) \vee \mathcal{X}_{V \setminus (A \cup B)}(t) \vee \mathcal{X}_V(t-k).$$

Since this relation holds for all $t \in \mathbb{Z}$, we have by (CI2) and (CI3)

$$\mathcal{X}_A(t) \perp\!\!\!\perp \mathcal{X}_B(t) \mid \mathcal{X}_{V \setminus (A \cup B)}(t) \vee \mathcal{X}_{A \cup B}(t-k-1),$$

which completes the induction step.

To show that (C.5) entails (C.4), we note that for $k \rightarrow \infty$ (C.5) yields

$$\mathcal{X}_A(t) \perp\!\!\!\perp \mathcal{X}_B(t) \mid \bigcap_{k>0} [\mathcal{X}_{V \setminus (A \cup B)}(t) \vee \mathcal{X}_{A \cup B}(t-k)]$$

for all $t \in \mathbb{Z}$. As in the proof of Proposition A.2, it follows that

$$\bigcap_{k>0} [\mathcal{X}_{V \setminus (A \cup B)}(t) \vee \mathcal{X}_{A \cup B}(t-k)] = \mathcal{X}_{V \setminus (A \cup B)}(t),$$

which concludes the proof of (C.4). $\square$

Proof of Theorem 4.3. Suppose that A , B , and S are disjoint subsets of V such that $A \rtimes_p B \mid S$. Let ξ be any $\mathcal{X}_A(\infty)$ measurable random variable with $\mathbb{E}|\xi| < \infty$, where $\mathcal{X}_A(\infty) = \vee_{t \in \mathbb{Z}} \mathcal{X}_A(t)$ denotes the σ -algebra generated by X_A . Then $\xi(t) = \mathbb{E}(\xi \mid \mathcal{X}_A(t))$ is a martingale and converges to ξ in L^1 as t tends to infinity. Thus, we obtain on the one hand, as $t \rightarrow \infty$,

$$\mathbb{E}(\xi(t) \mid \mathcal{X}_{S \cup B}(t)) \rightarrow \mathbb{E}(\xi \mid \mathcal{X}_{S \cup B}(\infty)) \quad \text{in } L^1. \quad (\text{C.9})$$

On the other hand, since $\xi(t) \perp\!\!\!\perp \mathcal{X}_B(t) \mid \mathcal{X}_S(t)$ by Lemma 4.2, we have, as $t \rightarrow \infty$,

$$\mathbb{E}(\xi(t) \mid \mathcal{X}_{S \cup B}(t)) = \mathbb{E}(\xi(t) \mid \mathcal{X}_S(t)) \rightarrow \mathbb{E}(\xi \mid \mathcal{X}_S(\infty)) \quad \text{in } L^1. \quad (\text{C.10})$$

Since the limits in (C.9) and (C.10) must be equal in L^1 and, thus, also almost surely, this proves that $\mathcal{X}_A(\infty) \perp\!\!\!\perp \mathcal{X}_B(\infty) \mid \mathcal{X}_S(\infty)$. $\square$




FIGURE C.2. (a) bi-directed path; (b) extended bi-directed path.

Proof of Theorem 4.6. For the proof of the first part of the global Granger-causal Markov property, let A and B subsets such that all B -pointing paths between A and B are p -blocked given $B \cup C$. We note that each B -pointing path π is of the form $\pi = \langle \tilde{\pi}, e \rangle$, where e is a directed edge $u \rightarrow b$ for some $b \in B$. Thus, π is p -blocked given $B \cup C$ if and only if $u \in B \cup C$ or $\tilde{\pi}$ is p -blocked given $B \cup C$. Therefore, if all B -pointing paths between A and B are p -blocked given $B \cup C$, then A and $\text{pa}(B) \setminus (B \cup C)$ are p -separated given $B \cup C$ and we obtain by Lemma 4.2

$$\mathcal{X}_{\text{pa}(B) \setminus (B \cup C)}(t) \perp\!\!\!\perp \mathcal{X}_A(t) \mid \mathcal{X}_{B \cup C}(t).$$

Since, in particular, every edge $a \rightarrow b$ for some $a \in A$ and $b \in B$ is p -connecting, it follows that A and $\text{pa}(B)$ are disjoint. Thus, we get by the block-recursive Granger-causal Markov property

$$\mathcal{X}_B(t+1) \perp\!\!\!\perp \mathcal{X}_A(t) \mid \mathcal{X}_{\text{pa}(B) \cup S \cup B}(t).$$

Applying the contraction property to this and the previous relation, we find that X_A is Granger-noncausal for X_B with respect to $\mathcal{X}_{A \cup B \cup S}$.

For the proof of the second part, let $U = A \cup B \cup S$ and assume that every extended bi-pointing path between A and B is p -blocked given U . Firstly, we note that a bi-directed path π between $a \in A$ and $b \in B$ is of the form $\pi = \langle e_1, \tilde{\pi}, e_n \rangle$, where e_1 and e_n are directed edges $a \leftarrow v_1$ and $v_{n-1} \rightarrow b$, respectively (Fig. C.2 a). Thus, π is p -blocked given U if and only if $v, w \in U$ or $\tilde{\pi}$ is p -blocked given U . This implies that, if all bi-directed paths between A and B are p -blocked given U , $\text{pa}(A) \setminus U$ and $\text{pa}(B) \setminus U$ are p -separated given U .

The argument can be adapted to the more general case of extended bi-directed paths: let S_A be the subset of $s \in S$ that are linked to A by an undirected p -connecting path and define S_B similarly. Then, intuitively, $\text{pa}(A \cup S_A) \setminus U$ and $\text{pa}(B \cup S_B) \setminus U$ should be p -separated given U (Fig. C.2 b).

More precisely, let $S_0 = \{s \in S \mid \text{pa}(s) \subseteq U\}$, which in particular includes all $s \in S$ that have no parents. Furthermore, let S_A be the set of all $s \in S \setminus S_0$ such that every extended bi-pointing path between s and B is p -blocked given U and set $S_B = S \setminus (S_0 \cup S_A)$. Notice that for all $s \in S_B$ there exists an extended bi-pointing path between s and B that is p -connecting given U . We show that every extended bi-pointing path between $A \cup S_A$ and $B \cup S_B$ is p -blocked given U . Since all extended bi-pointing paths between $A \cup S_A$ and B must be p -blocked by assumption on A and B or by definition of S_A , we only have to show that all ext-bi-pointing paths between $A \cup S_A$ and S_B are p -blocked given U . Suppose to the contrary that π is an extended bi-pointing path between $A \cup S_A$ and $s \in S_B$ that is p -connecting given U . Then, as mentioned above, there exists a p -connecting extended bi-pointing path π_s between s and B . If s is a p -collider on the composed extended bi-pointing path $\pi' = \langle \pi, \pi_s \rangle$ then π' is p -connecting given U contradicting the assumption

about A and B . Otherwise, if s is a p -noncollider, the two adjacent edges must be undirected (i.e. $— s —$) because extended bi-pointing paths never have a tail at either endpoint. Noting that there exists some $v \in \text{pa}(s) \setminus U$ since $s \notin S_0$, we set $\pi' = \langle \pi, s \leftarrow v \rightarrow s, \pi_s \rangle$. Then s is both times a p -collider on π' and, thus, π' is p -connecting given U . Since π' is also an extended bi-pointing path, this contradicts again the assumption about A and B .

Since in particular all bi-pointing paths between $A \cup S_A$ and $B \cup S_B$ are p -blocked given U , we have

$$\text{pa}(A \cup S_A) \setminus U \propto_p \text{pa}(A \cup S_B) \setminus U \mid U.$$

Thus, we obtain by Lemma 4.2

$$\mathcal{K}_{\text{pa}(A \cup S_A) \setminus U}(t) \perp\!\!\!\perp \mathcal{K}_{\text{pa}(B \cup S_B) \setminus U}(t) \mid \mathcal{K}_U(t). \quad (\text{C.11})$$

Since $\text{pa}(A \cup S_A) \subseteq V \setminus \text{pa}(B \cup S_B)$ and $\text{pa}(S_0) \subseteq U$, the block-recursive Granger-causal Markov property implies

$$\mathcal{K}_{B \cup S_B \cup S_0}(t+1) \perp\!\!\!\perp \mathcal{K}_{\text{pa}(A \cup S_A) \setminus U}(t) \mid \mathcal{K}_{U \cup \text{pa}(B \cup S_B)}(t) \quad (\text{C.12})$$

and further with (C.11)

$$\mathcal{K}_{B \cup S_B \cup S_0}(t+1) \perp\!\!\!\perp \mathcal{K}_{\text{pa}(A \cup S_A) \setminus U}(t) \mid \mathcal{K}_U(t). \quad (\text{C.13})$$

Moreover, since undirected paths are special cases of extended bi-pointing paths, we find that every undirected path between $A \cup S_A$ and $B \cup S_B$ intersects S_0 . Then, by a standard argument of graph theory (e.g., Whittaker 1990, Lemma 3.3.3), there exists a partition (A^*, B^*, S_0) of V such that $A \cup S_A \subseteq A^*$, $B \cup S_B \subseteq B^*$, and every undirected path between A^* and B^* intersects S_0 ; in particular, this implies $\text{ne}(A \cup S_A) \subseteq S_0$. Thus, we obtain by the block-recursive Granger-causal Markov property

$$\mathcal{K}_{A \cup S_A}(t+1) \perp\!\!\!\perp \mathcal{K}_{B \cup S_B}(t+1) \mid \mathcal{K}_V(t) \vee \mathcal{K}_{S_0}(t+1).$$

Together with

$$\mathcal{K}_{A \cup S_A \cup S_0}(t+1) \perp\!\!\!\perp \mathcal{K}_{V \setminus (U \cup \text{pa}(A \cup S_A))}(t) \mid \mathcal{K}_{U \cup \text{pa}(A \cup S_A)}(t),$$

which also follows from the block-recursive Granger-causal Markov property, this implies

$$\mathcal{K}_{A \cup S_A}(t+1) \perp\!\!\!\perp \mathcal{K}_{B \cup S_B}(t+1) \mid \mathcal{K}_{U \cup \text{pa}(A \cup S_A)}(t), X_{S_0}(t+1). \quad (\text{C.14})$$

Applying (C.14) to (C.13) and (C.14), we finally obtain

$$\mathcal{K}_{A \cup S_A}(t+1) \perp\!\!\!\perp \mathcal{K}_{B \cup S_B}(t+1) \mid \mathcal{K}_U(t) \vee \mathcal{K}_{S_0}(t+1),$$

from which the desired relation follows by (C.12).

Finally, to see that (GC) entails (BC), let $S = \text{pa}(B)$ and $A = V \setminus S$ for an arbitrary subset B of V . Then the first relation in (BC) follows directly from the global Granger-causal Markov property. The second relation in (BC) can be derived similarly. $\square$

Proof of Corollary 4.7. Suppose that all paths between A and B are p -blocked given S . We show that then all B -pointing paths between A and B are p -blocked given $S \cup B$, which implies by the global Granger-causal Markov property that X_A is Granger-noncausal for X_B with respect to $\mathcal{K}_{A \cup B \cup S}$.

We firstly note that, in particular, every B -pointing path π between A and B are p -blocked given S and, if π does not contain any intermediate nodes in B , also

p -blocked given $S \cup B$. Now, suppose that π is a B -pointing path between A and B with some intermediate nodes in B . Then π can be partitioned as $\pi = \langle \pi_1, \pi_2 \rangle$ where π_1 is a path between A and some $b \in B$ with no intermediate nodes in B . Because of the assumption, the path π_1 is p -blocked given S and, since it has no intermediate vertices in B , also given $S \cup B$. It follows that all B -pointing paths between A and B are p -blocked given $S \cup B$.

The other two cases $X_B \nrightarrow X_A$ [$\mathcal{X}_{A \cup B \cup S}$] and $X_A \approx X_B$ [$\mathcal{X}_{A \cup B \cup S}$] can be derived similarly. $\square$

APPENDIX D. P-SEPARATION IN MIXED GRAPHS

The definition of p -separation presented in this paper is based on paths that may be self-intersecting. This leads to simpler conditions than in the original definition by Levitz et al. (2001). The latter is formulated in terms of paths on which all intermediate vertices are distinct, that is, these paths are not self-intersecting; such paths are called trails. According to Levitz et al. (2001), a trail between vertices a and b is said to be p -active relative to S if

- (i) every p -collider (head-no-tail node) on π is in $\text{an}(S)$, and
- (ii) every p -noncollider v is either not in S or it has two adjacent undirected edges $(-v-)$ and $\text{pa}(v) \setminus S \neq \emptyset$.

Otherwise the trail is p -blocked relative to S . Let A , B , and S be disjoint subsets of V . Then S p -separates A and B if all trails between A and B are p -blocked relative to S .

The following proposition shows that the two notions of p -separation are equivalent.

Proposition D.1. *Let $G = (V, E)$ be a mixed graph and A , B , S disjoint subsets of V . Then there exists a p -active trail between A and B relative to S if and only if there exists a p -connecting path between A and B given S .*

Proof. Suppose that π is a trail between two vertices a and b that is p -active relative to S . If all p -colliders on π are in S and all p -noncolliders are outside S , then π is also p -connecting given S . Otherwise, π is p -blocked by vertices $u_{j_1}, \dots, u_{j_r}$ on the path. If u_{j_i} is a p -collider then $u_{j_i} \in \text{an}(S)$ since π is p -active. Hence there exists a directed path $\tau_i = \langle u_{j_i} \rightarrow \dots \rightarrow s_i \rangle$ for some $s_i \in S$ such that all intermediate vertices on τ_i are not in S and we set $\sigma_i = \langle \tau_i, \bar{\tau}_i \rangle$. On the other hand, if u_{j_i} is a p -noncollider on π , then the two edges adjacent to u_{j_i} are undirected. Thus, there exists $w_i \in \text{pa}(u_{j_i}) \setminus S$ and we set $\sigma_i = \langle u_{j_i} \leftarrow w_i \rightarrow u_{j_i} \rangle$. Now, let π_i be the subpath of π between $u_{j_{i-1}}$ and u_{j_i} with $u_{j_0} = a$ and $u_{j_{r+1}} = b$ and set

$$\pi' = \langle \pi_0, \sigma_1, \pi_1, \sigma_2, \dots, \pi_{r-1}, \sigma_r, \pi_r \rangle.$$

Then all p -colliders on π' are in S and all p -noncolliders are not in S , which yields that π' is p -connecting given S .

Conversely, suppose that π is a p -connecting path between a and b given S . Let u_{j_1} be the first vertex on the path that occurs more than once. Then π can be partitioned as $\pi = \langle \pi'_0, \lambda_1, \pi_1 \rangle$ such that u_{j_1} is an endpoint, but not an intermediate node of π'_0 and π_1 . Noting that π'_0 is already a trail, we continue to partition π_1 in

the same way. After finitely many steps, we obtain the partition

$$\pi = \langle \pi'_0, \lambda_1, \pi'_1, \lambda_2, \dots, \pi'_{r-1}, \lambda_r, \pi'_r \rangle$$

such that the subpaths π'_j are all trails. Thus, the shortened path $\pi' = \langle \pi'_0, \dots, \pi'_r \rangle$ is also a trail. We show that π' is a p -active trail relative to S . We firstly note that all subtrails π'_j are p -connecting and hence p -active. We therefore have to show that the vertices u_{j_i} satisfy the conditions for a p -active trail.

Suppose that u_{j_i} is a p -collider that is not in S . Then at least one of the edges adjacent to u_{j_i} has an arrowhead at u_{j_i} and we may assume that π'_{i-1} is u_{j_i} -pointing (otherwise consider the reverse path). Since $u_{j_i} \notin S$, it must be a p -noncollider on π and hence λ_i starts with a tail at u_{j_i} . On the other hand, since u_{j_i} must be a p -noncollider on all its occurrences on π and π'_i does not start with a tail, the loop λ_i cannot be a directed path (otherwise u_{j_i} would not be a p -collider on $\langle \lambda_i, \pi'_i \rangle$). Consequently there exists an intermediate vertex w_i such that the subpath between u_{j_i} and w_i is directed and w_i is a p -collider. It follows that $w_i \in S$ and $u_{j_i} \in \text{an}(S)$.

Next, suppose that u_{j_i} is a p -noncollider on π' that is in S . Since u_{j_i} has been a p -collider on π , the two edges adjacent to u_{j_i} on π' must be undirected and λ_i must be a bi-directed path. Hence λ_i is of the form $\lambda_i = \langle u_{j_i} \leftarrow w_i, \lambda'_i \rangle$ with $w_i \notin S$ (since w_i is a p -noncollider and π is p -connecting). Therefore, the set $\text{pa}(u_{j_i}) \setminus S$ is not empty and u_{j_i} satisfies the above condition (ii). Altogether it follows that π' is p -active relative to S . $\square$

In a remark on our simplified version of p -separation, Levitz et al. (2001) argue that there are infinitely many possibly self-intersecting paths in a graph as opposed to finitely many trails. The following lemma shows that it is possible to restrict the search for p -connecting paths in G to a finite number of paths, namely all paths in which no edge occurs twice with the same orientation.

Lemma D.2. *Let $G = (V, E)$ be a mixed graph and suppose that π is a p -connecting path of the form $\pi = \langle \pi_1, e, \pi_2, e, \pi_3 \rangle$, where e is an oriented edge between some vertices u and v . Then the shortened path $\pi' = \langle \pi_1, e, \pi_3 \rangle$ is also p -connecting.*

Proof. Since π is p -connecting, the two subpaths $\langle \pi_1, e \rangle$ and $\langle e, \pi_3 \rangle$ are p -connecting. This implies that also π' is p -connecting as every intermediate vertex has the same p -collider/noncollider status as in the corresponding subpath. $\square$

REFERENCES

- Andersson, S. A., Madigan, D. and Perlman, M. D. (2001). Alternative Markov properties for chain graphs. *Scandinavian Journal of Statistics* **28**, 33–85.
- Baccalá, L. A. and Sameshima, K. (2001). Partial directed coherence: a new concept in neural structure determination. *Biological Cybernetics* **84**, 463–474.
- Boudjellaba, H., Dufour, J. M. and Roy, R. (1992). Testing causality between two vectors in multivariate autoregressive moving average models. *JASA* **87**, 1082–1090.
- Brillinger, D. R. (1988a). Maximum likelihood analysis of spike trains of interacting nerve cells. *Biological Cybernetics* **59**, 189–200.
- Brillinger, D. R. (1988b). The maximum likelihood approach to the identification of neuronal firing systems. *Annals of Biomedical Engineering* **16**, 3–16.
- Brillinger, D. R. (1996). Remarks concerning graphical models for time series and point processes. *Revista de Econometria* **16**, 1–23.

- Cox, D. R. and Wermuth, N. (1996). *Multivariate Dependencies - Models, Analysis and Interpretation*. Chapman & Hall, London.
- Dahlhaus, R. (2000). Graphical interaction models for multivariate time series. *Metrika* **51**, 157–172.
- Dahlhaus, R. and Eichler, M. (2003). Causality and graphical models in time series analysis. In P. Green, N. Hjort and S. Richardson (eds), *Highly structured stochastic systems*, University Press, Oxford.
- Dawid, A. P. (1979). Conditional independence in statistical theory (with discussion). *Journal of the Royal Statistical Society Series B* **41**, 1–31.
- Edwards, D. (2000). *Introduction to Graphical Modelling*. 2nd edn, Springer, New York.
- Eichler, M. (2005a). A graphical approach for evaluating effective connectivity in neural systems. *Philosophical Transactions of The Royal Society B* **360**, 953–967.
- Eichler, M. (2005b). Graphical modelling of multivariate time series with latent variables. *Technical report*, University of Heidelberg.
- Eichler, M. (2006a). Granger causality and path diagrams for multivariate time series. *Journal of Econometrics* (in press).
- Eichler, M. (2006b). Graphical modelling of dynamic relationships in multivariate time series. In M. Winterhalder, B. Schelter and J. Timmer (eds), *Handbook of Time Series Analysis*, Wiley-VCH.
- Fan, J. and Yao, Q. (2003). *Nonlinear Time Series: Nonparametric and Parametric Methods*. Springer, New York.
- Florens, J. P. and Mouchart, M. (1982). A note on noncausality. *Econometrica* **50**, 583–591.
- Florens, J. P., Mouchart, M. and Rolin, J. M. (1990). *Elements of Bayesian Statistics*. Marcel Dekker, New York.
- Frydenberg, M. (1990). The chain graph Markov property. *Scandinavian Journal of Statistics* **17**, 333–353.
- Granger, C. W. J. (1969). Investigating causal relations by econometric models and cross-spectral methods. *Econometrica* **37**, 424–438.
- Hsiao, C. (1982). Autoregressive modeling and causal ordering of econometric variables. *Journal of Economic Dynamics and Control* **4**, 243–259.
- Koster, J. T. A. (1999). On the validity of the Markov interpretation of path diagrams of Gaussian structural equations systems with correlated errors. *Scandinavian Journal of Statistics* **26**, 413–431.
- Lauritzen, S. L. (1996). *Graphical Models*. Oxford University Press, Oxford.
- Levitz, M., Perlman, M. D. and Madigan, D. (2001). Separation and completeness properties for AMP chain graph Markov models. *Annals of Statistics* **29**, 1751–1784.
- Lynggaard, H. and Walther, K. H. (1993). *Dynamic modelling with mixed graphical association models*. Master's thesis, Aalborg University.
- Moneta, A. and Spirtes, P. (2005). Graph-based search procedure for vector autoregressive models. *LEM Working Paper 2005/14*, Sant'Anna School of Advanced Studies, Pisa.
- Oxley, L., Reale, M. and Tunnicliffe Wilson, G. (2004). Finding directed acyclic graphs for vector autoregressions. In J. Antoch (ed.), *Proceedings in Computational Statistics 2004*, Physica-Verlag, Heidelberg, pp. 1621–1628.
- Pearl, J. (1988). *Probabilistic Inference in Intelligent Systems*. Morgan Kaufmann, San Mateo, California.
- Pearl, J. and Paz, A. (1987). Graphoids: a graph based logic for reasoning about relevancy relations. In B. D. Boulay, D. Hogg and L. Steel (eds), *Advances of Artificial Intelligence-II*, North-Holland, Amsterdam, pp. 357–363.

- Reale, M. and Tunnicliffe Wilson, G. (2001). Identification of vector AR models with recursive structural errors using conditional independence graphs. *Statistical Methods and Applications* **10**, 49–65.
- Rothman, P. (ed.) (1999). *Nonlinear Time Series Analysis of Economic and Financial Data*. Kluwer Academic Publishers.
- Stanghellini, E. and Whittaker, J. (1999). Analysis of multivariate time series via a hidden graphical model. *Proceedings of the 7th International Workshop on Artificial Intelligence and Statistics*, Morgan Kaufmann, San Mateo, CA, pp. 250–254.
- Talih, M. and Hengartner, N. (2005). Structural learning with time-varying components: tracking the cross-section of financial time series. *Journal of the Royal Statistical Society Series B* **67**, 321–341.
- Tong, H. (1993). *Non-Linear Time Series: A Dynamical System Approach*. Oxford University Press, Oxford.
- Veijanen, A. (1990). On conditionally mixing processes. *Statistics and Probability Letters* **9**, 59–65.
- Whittaker, J. (1990). *Graphical Models in Applied Multivariate Statistics*. John Wiley, Chichester.